ACCEPTED

JUN 15 1990

Under the Federal Insecticide, Fungicues, and Rodenticide Act, as amended, for the pest cide rightness under EPA Reg. No. 202-526

LILLY/MILLER HOSE 'N GO WEED & FEED

15-0-0

Contains TRIMEC

Control weeds and feed grass in one application. Just attach to a garden hose and spray.

Kills dandelions, plantains, clover, chickweed, spurge, oxalis and many other lawn weeds.

Ready to use. Covers up to 3,750 sq. ft.

902-435 Label

٠..

ACTIVE INGREDIENTS:

*Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid 5.673% **Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy)

propionic acid 2.738%

***Dimethylamine Salt of Dicamba

(3,6-dichloro-o-anisic acid) 0.629% INERT INGREDIENTS 90.96%

INERT INGREDIENTS 90.96%

*2,4-Dichlorophenoxyacetic acid equivalent 4.708% (0.44 lb. per gallon). Isomer specific by AOAC method.

**2-(2-methyl-4-chlorophenoxy)propionic acid equivalent 2.272% (0.21 lb. per gallon)

***Dicamba (3,6-dichloro-o-anisic acid) equivalent 0.522% (0.048 lb. per gallon)

TRIMEC is the Reg. T.M. of PBI/Gordon Corporation.

Keep out of reach of children CAUTION

See back panel for additional precautionary statements.

NET CONTENTS 24 FL. OZ.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin or clothing. When handling this product, wear chemical resistant gloves. Wear goggles, long pants, long-sleeved shirt and shoes when applying this product. In case of contact, immediately flush eyes or skin with plenty of water. Avoid contamination of food or feedstuffs. Do not reenter or permit others to reenter treated areas until sprays have dried.

STATEMENT OF PRACTICAL TREATMENT: In case of contact, wash skin with

BEST AVAILABLE COPY

302. 49

rupel

plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention.

ENVIRONMENTAL HAZARDS: This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget / plants. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwaters.

OBSERVE ALL PRECAUTIONS AND FOLLOW READ THE ENTIRE LABEL FIRST. STOP. DIRECTIONS CAREFULLY.

DIRECTIONS FOR USE It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

WHEN TO USE: Spray at any time during the growing season when weeds are actively growing. On new lawns, wait until the grass has been mowed two or three times and has reached a height of 2 inches before making an overall spray treatment. Wait 3 weeks before putting new grass seed or , stolons down in the bare spots. Poor weed control may result if spray is applied during drought or just before rain. For best results, mow 1 to 2 days before treating. Do not water lawn within 24 hours after treatment.

WHERE TO USE: HOSE 'N GO WEED & FEED may be used on Bahia, Bentgrass*, Bermuda, Bluegrass, Fescue, Rye or Zoysia lawns to control broadleaf HOSE 'N GO WEED & FEED controls Bedstraw, Buckhorn, Buttercup, Chickweed, Clover, Dandelion, Dock, English Daisy, Ground Ivy, Henbit, All of Japanese Clover (Black Medic), Knotweed, Lambsquarter, Mallow, Morningglory, Oxalis, Pigweed, Plantain, Poison Ivy, Poison Oak, Purslane, Sheep Sorrel, Shepardspurse, Speedwell, Spurge, Yarrow, and most other broadleaved lawn weeds.

*See Use Precautions.

HOW TO USE: Shake well before using. 1. Connect spray nozzle to garden hose. Be sure that valve is in "off" position. 2. Remove bottle cap and screw bottle onto nozzle. Turn on water. to "on" position to start flow of water. Use valve to shut off flow of water at any time. Spray directly over area requiring application (Check handy measuring guide to determine: " evenly and uniformly. remaining area to be covered).

USE PRECAUTIONS: Do not allow spray to drift to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants since injury may result. Do not spray where grass is emerging from dormancy. Do not apply to newly seeded lawn grasses until well established. Do not s, 'ay', when the temperature exceeds 85 degrees F. Seed can be safely sown 3 weeks after application. Do not use on carpetgrass, dichondra, Gt.

BEST AVAILABLE COPY

Augustine, Centipede, closely-mowed bentgrass, or on lawns or turf where - desirable clovers are present. Do not apply to lawn areas above the root zone of trees and ornamentals, or within the dripline.

GUARANTEED FERTILIZER ANALYSIS
Total Nitrogen (N) . . . 15% /
15% Urea Nitrogen /
Primary nutrients from Urea.

STORAGE AND DISPOSAL

Keep from freezing. Store this product in its original container out of reach of children and domestic animals. Do not reuse container. Rinse throughly before discarding and securely wrap original container in several layers of newspaper and discard in trash.

IOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

THE CHAS. H. LILLY CO. PORTLAND, OREGON 97218 / EPA Reg. No. 802-586 EPA Est. 802-OR-1 91111