

The  
Chas. H.  
LILLY Co.

LILLY/MILLER.  
*Cooke*

PM 23 802-586 1044  
7737 N.E. KILLINGSWORTH • PORTLAND, OREGON 97218 • (503) 256-4600  
• FAX (503) 255-7552

February 27, 1993

I, being an authorized representative of The Chas. H. LILLY Co., certify that all containers of Lilly/Miller Hose 'n Go Weed & Feed, EPA Reg. No. 802-586 produced by June 15, 1994 will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing-use products. I further certify that all containers of said product sold or distributed by this company by January 1, 1995 will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing-use products.

*Molly Klettke*

Molly Klettke, Registration Specialist

204

LILLY/MILLER  
HOSE 'N GO WEED & FEED 15-0-0

Contains TRIMEC

Control weeds and feed grass in one application. Just attach to a garden hose and spray.

Ready to use. Covers up to 3,750 sq. ft.

ACTIVE INGREDIENTS:

*Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid .....	5.673%
**Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid .....	2.738%
***Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid) .....	0.629%
) INERT INGREDIENTS .....	90.96%

\*2,4-Dichlorophenoxyacetic acid equivalent 4.708% (0.44 lb. per gallon). Isomer specific by AOAC method.

\*\*2-(2-methyl-4-chlorophenoxy)propionic acid equivalent 2.272% (0.21 lb. per gallon)

\*\*\*Dicamba (3,6-dichloro-o-anisic acid) equivalent 0.522% (0.048 lb. per gallon)

TRIMEC is the Reg. T.M. of PBI/Gordon Corporation.

Keep out of reach of children

CAUTION

See back panel for additional precautionary statements.

NET CONTENTS 24 FL. OZ.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin or clothing. When using this product, wear goggles, long-sleeved shirt, long pants, socks, shoes and chemical resistant gloves. In case of contact, immediately flush eyes or skin with plenty of water. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. remove saturated clothing as soon as possible and shower. Avoid contamination of food or feedstuffs. Do not allow people (other than applicator) or pets on treatment area during application. Do not reenter or permit others to reenter treated areas until sprays have dried.

STATEMENT OF PRACTICAL TREATMENT: In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and

get medical attention.

**ENVIRONMENTAL HAZARDS:** This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwaters.

**STOP. READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. The maximum number of broadcast applications per treatment site is 2 per year.

**WHEN TO USE:** Spray at any time during the growing season when weeds are actively growing. On new lawns, wait until the grass has been mowed two or three times and has reached a height of 2 inches before making an overall spray treatment. Wait 3 weeks before putting new grass seed or stolons down in the bare spots. Poor weed control may result if spray is applied during drought or just before rain. For best results, mow 1 to 2 days before treating. Do not water lawn within 24 hours after treatment.

**WHERE TO USE:** HOSE 'N GO WEED & FEED may be used on Bahia, Bentgrass\*, Bermuda, Bluegrass, Fescue, Rye or Zoysia lawns to control broadleaf weeds. HOSE 'N GO WEED & FEED controls Bedstraw, Buckhorn, Buttercup, Chickweed, Clover, Dandelion, Dock, English Daisy, Ground Ivy, Henbit, Japanese Clover (Black Medic), Knotweed, Lambsquarter, Mallow, Morningglory, Oxalis, Pigweed, Plantain, Poison Ivy, Poison Oak, Purslane, Sheep Sorrel, Shepardspurse, Speedwell, Spurge, Yarrow, and most other broadleaved lawn weeds.

\*See Use Precautions.

**HOW TO USE:** Shake well before using. 1. Connect spray nozzle to garden hose. Be sure that valve is in "off" position. 2. Remove bottle cap and screw bottle onto nozzle. Turn on water. 3. Switch valve to "on" position to start flow of water. Use valve to shut off flow of water at any time. Spray directly over area requiring application evenly and uniformly. (Check handy measuring guide to determine remaining area to be covered).

**USE PRECAUTIONS:** Do not allow spray to drift to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants since injury may result. Do not spray where grass is emerging from dormancy. Do not apply to newly seeded lawn grasses until well established. Do not spray when the temperature exceeds 85 degrees F. Seed can be safely sown 3 weeks after application. Do not use on carpetgrass, dichondra, St.

4044

Augustine, Centipede, closely-mowed bentgrass, or on lawns or turf where desirable clovers are present. Do not apply to lawn areas above the root zone of trees and ornamentals, or within the dripline.

**GUARANTEED FERTILIZER ANALYSIS**

Total Nitrogen (N) . . . . 15%  
15% Urea Nitrogen  
Primary nutrients from Urea.

**STORAGE AND DISPOSAL**

Keep from freezing. Store this product in its original container out of reach of children and domestic animals. Do not reuse container. Rinse thoroughly before discarding and securely wrap original container in several layers of newspaper and discard in trash.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

THE CHAS. H. LILLY CO.  
PORTLAND, OREGON 97218  
EPA Reg. No. 802-586  
EPA Est. 802-OR-1  
91111

