

MINIMAX

MALATHION

INSECTICIDE



AMERICAN
DISINFECTANT COMPANY

CAUTION

Keep Out of Reach of Children

See container for additional caution statements.

1.000ml liquid

DIRECTIONS

This product is for use in Homes, Stores, Motels, Hotels, Hospitals, Office Buildings, Apartment Buildings and Warehouses.

For Fleas: A thorough wetting of all cracks and crevices and treatment of entire floor surfaces is necessary. Special attention should be paid to area where animals sleep. (DO NOT SPRAY ANIMALS). Keep children and animals out of area treated for 4 hours after treatment. Repeat treatment if reinfestation occurs.

For Bedbugs: Apply the liquid insecticide to the entire surface of the frame of the bed, the springs, and lightly to all surfaces of the mattress. Apply generously to the walls adjacent to the furniture in such a manner as to cover all surfaces over which the insects would travel in order to reach their host. Repeat treatment if reinfestation occurs.

CAUTION

May be absorbed through skin. Avoid breathing of spray mist and avoid skin contact. If contact is made with skin, immediately wash thoroughly with warm water and soap. For eyes, flush with water for several minutes and get prompt medical attention. Harmful if swallowed. If taken internally, administer an emetic and call a physician. Keep out of reach of children. Do not use on plants or animals. Combustible mixture: Do not use, pour, spill or store near heat or open flame or in such a manner as to contaminate feed or foodstuffs. Do not smoke while spraying.

Do not reuse container. Destroy when empty.

Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

MALATHION, upon repeated or prolonged use, may cause cholinesterase inhibition. If poisoning occurs, get prompt medical attention.

ACCEPTED
July 19, 1971
809-11
with comments

ACCEPTED WITH COMMENTS