

50 LBS. NET
AGRIC. CHEM. CO. OF AMERICA

10-0-0

PLANT NUTRITION DIVISION, AGRIC. RESEARCH SERVICE

GUARANTEED ANALYSIS

8-27-2003

Total Nitrogen (N)	10.00%
1.33 Water Insoluble Nitrogen	
Available Phosphoric Acid (P ₂ O ₅)	6.00%
Soluble Potash (K ₂ O)	4.00%
Potential Acidity 500 Ppb.	
Calcium Carbonate Equivalent for 100.	

*Reads also as "Minimum Guaranteed Analysis"

HERBICIDE INGREDIENT STATEMENT

ACTIVE INGREDIENTS BY WEIGHT:

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	0.55%
Equivalent to 0.46% 2,4-dichlorophenoxyacetic acid	
Dimethylamine salt of "Silvex" 2-(2,4,5-trichlorophenoxy) propionic acid	0.20%
Equivalent to 0.17% "Silvex" 2-(2,4,5-trichlorophenoxy) propionic acid	
Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	0.28%
Equivalent to 0.23% 2-(2-methyl-4-chlorophenoxy) propionic acid	

INERT INGREDIENTS

(Relative to herbicide content)

98.97%

USDA REG. NO. 827-203

Controls: Sanderson, Plantain, Buckhorn, Shepherd's Purse, Chickweed, Stitchwort, Knotweed, Clover, Sheep Sorrel and Henbit in lawns, golf course fairways, aprons, tees, roughs and other turf (Not pasture) areas.

INSTRUCTIONS

1. Apply uniformly at the rate of 4.0 lbs. per acre with a fertilizer spreader. This will give the recommended herbicide treatment.

SUGGESTED APPLICATIONS

Apply to:	Rate	Comments
Jack-o'-lantern	2.0/1.0	Use special spreader
Lawn Care	4.0/1.0	

2. For best results, apply during the dormant or actively growing period.

3. Increased effectiveness is obtained by applying when the grass is moist such as early in the morning after a dew or follow a shower. Avoid, if possible, applying ahead of a heavy rain.
4. Do not apply under the drip line of trees or shrubs especially dogwood, willow and locust trees. Avoid drift or dust to susceptible plants. Be careful that water from treated areas does not accumulate around trees or shrubs.
5. May cause temporary injury to bent grass and clover.
6. May cause injury to grass seedlings on newly seeded ground.
7. Follow the Caution Statement.

-CAUTION-
KEEP OFF OF REACH OF CHILDREN

HARMFUL IF SWALLOWED: May cause irritation of eyes, nose, throat, and skin. Avoid inhalation of dust. Avoid contact with skin, eyes and clothing. In case of contact with skin or eyes, flush with plenty of water. For eyes get medical attention. Wash thoroughly with soap and water after handling this material and before eating or smoking.

Keep away from animals. Do not contaminate water used for irrigation or domestic purposes. Do not apply on desired flowers, shrubs, fruits, or vegetables. Do not store near and avoid contamination of foods, seeds, fungicides or fertilizers. Exercise care when disposing of first lawn clippings. Wash spreader after use.

Use this product only for recommended purposes and at recommended dosages.

WARRANTY

Seller makes no warranty, express or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risks to persons or property arising out of the improper use or careless handling of this material and accepts the product on these conditions.

MANUFACTURED BY
AGRI-CHEMICAL COMPANY
DIVISION OF CHEMICAL COMPANY
MEMPHIS, TENNESSEE

50 LBS. NET
AGRICO COUNTRY CLUB F

10-6-4

25% OF NITROGEN CONTENT FROM

GUARANTEED FERTILIZER

Total Nitrogen (N)
1.5% Water Insoluble Nitrogen

Available Phosphoric Acid (P₂O₅)

Soluble Potash (K₂O)
Potential Acidity 500 Lbs.
Calcium Carbonate Equivalent Per

*Reads also as "Minimum Guaranteed Analy

HERBICIDE INGREDIENT

ACTIVE INGREDIENTS BY WEIGHT:

Dimethylamine salt of 2,4-dichloro
acid

Equivalent to 0.46% 2,4-dichlorop
acid

Dimethylamine salt of "Silvex" 2-
phenoxy) propionic acid

Equivalent to 0.17% "Silvex" 2-(2-
phenoxy) propionic acid

Dimethylamine salt of 2-(2-methyl)
propionic acid

Equivalent to 0.23% 2-(2-methyl-4
propionic acid

INERT INGREDIENTS

(Relative to Herbicide Content)

Controls Dandelion, Plantain, Buck
Chickweed, Stitchwort, Knotweed,
Henbit in lawns, golf course fair
and other turf (Not pasture) areas

DIRECTIONS

1. Apply uniformly at the rate of 400 lb per acre with a broadcast spreader. This rate gives the recommended control.

SUGGESTED SPREADER

Agri-Spred	6-1/2	Sc
Jackson	6-1/2	Jet
Lawn Beauty	6-1/2	

2. For best results, apply when the weeds are young in the Fall or Spring seasons.

3. Increased effectiveness is obtained by applying when the grass is moist such as early in the morning after a dew or following a shower. Avoid, if possible, applying ahead of a heavy rain.
4. Do not apply under the dripline of trees or shrubs especially dogwood, willow and locust trees. Avoid drift of dust to susceptible plants. Be careful that water from treated areas does not accumulate around trees or shrubs.
5. May cause temporary injury to Bent grass and clover.
6. May cause injury to grass seedlings on newly seeded ground.
7. Follow the Caution Statement.

CAUTION
KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED. May cause irritation of eyes, nose, throat, and skin. Avoid inhalation of dust. Avoid contact with skin, eyes and clothing. In case of contact with skin or eyes, flush with plenty of water. For eyes get medical attention. Wash thoroughly with soap and water after handling this material and before eating or smoking.

Keep away from animals. Do not contaminate water used for irrigation or domestic purposes. Do not apply on desired flowers, shrubs, fruits, or vegetables. Do not store near and avoid contamination of foods, seeds, fungicides or fertilizers. Exercise care when disposing of first lawn clippings. Wash spreader after use.

Use this product only for recommended purposes and at recommended dosages.

WARRANTY

Seller makes no warranty, express or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risks to persons or property arising out of the improper use or careless handling of this material and accepts the product on these conditions.

MANUFACTURED BY
AGRICO CHEMICAL COMPANY
DIVISION OF CONTINENTAL OIL COMPANY
MEMPHIS, TENNESSEE