


Pro-forma
Bag Copy - Lawn Fungicide



March 23, 1967

FRONT:

Net weight 25 lbs.
Covers 2500 sq. ft.
when used as directed.

Retail Price
\$6.95 per bag

Agrico
Lawn
Fungicide

Ingredient Statement

ACTIVE INGREDIENT:

2,4-Dichloro-6-o-chloranilino-
s-triazine (Common name is Dyrene)

2,4-Dichloro-6- <u>o</u> -chloranilino- s-triazine (Common name is Dyrene)	3%
INERT INGREDIENTS:	97%
	<u>100%</u>

U. S. Patent No. 2,720,480 and
Canadian Patent No. 546,027
Effective for purpose claimed.

Reg. U.S. and Canadian Patent Offices
by Chemagro Corporation U.S.D.A. Reg. No. 827-

(Note: Underlined letters in chemical nomenclature to be
italicized - not underlined.) Do not include in bag
copy.

CAUTION

KEEP OUT OF REACH OF CHILDREN

May be harmful if swallowed or inhaled. Avoid contact with eyes,
skin, or clothing. Avoid breathing of dust or spray mist. Wash
thoroughly with soap and warm water after handling. Wash con-
taminated clothing with soap and hot water before re-use. Do
not contaminate feed or food.

BACK:

Agrico
Lawn Fungicide
25 lbs. net weight

DIRECTIONS FOR USE

Agrico Lawn Fungicide effectively protects grasses from various
destructive diseases. Uniform distribution of granular material
is necessary for maximum control. For a preventive program,
begin applications before diseases appear and repeat treatments
as needed, usually at intervals of 10 to 14 days, except during
extended periods of cool, moist conditions when applications
every 7 days may be advisable. Agrico Lawn Fungicide applications
should be continued through the spring and summer months.

For control of lawn diseases, Brown Patch, Copper Spot, Dollar Spot, Leaf Spot, Melting Out and Snow Mold apply 6 to 10 lbs. of Agrico Lawn Fungicide per 1000 sq. feet. Repeat at 1 to 2 week intervals throughout the spring and summer months. Use recommended spreader settings for 10 lbs. per sq. ft. Spreader adjustments may be necessary to reduce quantities and to allow for walking speed and topography. Spreader settings are given as a guide only.

Suggested Spreader Settings

	2500 sq. ft. coverage		2500 sq. ft. coverage
Agri-Spred	8	Modern	8
Lawn Beauty	8½	Sears	8
Scott	7½	Jackson	8

WARRANTY

Agrico Chemical Company makes no warranty, expressed or implied, except that this product conforms to the chemical description on the label. Said company shall not be held responsible in any manner for any personal injury or property damage or other type of loss resulting from the handling, storage or use of this product. The buyer assumes all risk and liability therefrom and accepts and uses this product on these conditions.

Agrico Chemical Company
Memphis, Tennessee 38101

GUSSETS:


Agrico Lawn Fungicide

PM31 827-209

Front Cover - Lawn Fungicide

Back Cover

Net Weight 25 lbs.
Government of Canada
which must be directed to



March 25, 1967

Nettoli Price
\$6.95 per bag

Agrico
Lawn
Fungicide

Ingredients

2,4-Dichloro-6-g-chloranilino- g-triazine (Common name is Pyricne)	30
INERT INGREDIENTS	97
	<u>100</u>

U.S. Patent No. 2,720,450 and
Canadian Patent No. 546,027
Effective for purposes claimed.

Reg. U.S. and Canadian Patent Offices
by Chemagro Corporation U.S.D.A. Reg. No. 927-

(Note: Underlined letters in chemical nomenclature to be
included - not underlined.) Do not include in bag
copy.

CAUTION

KEEP OUT OF REACH OF CHILDREN

May be harmful if swallowed or inhaled. Avoid contact with eyes,
skin, or clothing. Avoid breathing of dust or spray mist. Wash
thoroughly with soap and warm water after handling. Wash con-
taminated clothing with soap and hot water before re-use. Do
not contaminate feed or food.

BACK:

Agrico
Lawn Fungicide
25 lbs. net weight



DIRECTIONS FOR USE

Agrico Lawn Fungicide effectively protects grasses from various
destructive diseases. Uniform distribution of granular material
is necessary for maximum control. For a preventive program,
begin applications before diseases appear and repeat treatments
as needed, usually at intervals of 10 to 14 days, except during
extended periods of cool, moist conditions when applications
every 7 days may be advisable. Agrico Lawn Fungicide applications
should be continued through the spring and summer months.

For control of lawn diseases, Brown Patch, Copper Spot, Dollar Spot, Leaf Spot, Melting, Cut and Grow Field apply 6 to 10 lbs. of Agrico Lawn Fungicide per 100 sq. ft. Repeat at 1 to 2 week intervals throughout the spring and summer months. Use recommended spreader settings for 10 lbs. per sq. ft. Spreader adjustment may be necessary to reduce quantities and to allow for walking speed and topography. Spreader settings are given as a guide only.

Suggested Spreader Settings

	2500 sq. ft. coverage		2500 sq. ft. coverage
Agrico-Spread	8	Modern	8
Lawn Beauty	8½	Scars	8
Scott	7½	Jackson	8

WARRANTY

Agrico Chemical Company makes no warranty, expressed or implied, except that this product conforms to the chemical description on the label. Said company shall not be held responsible in any manner for any personal injury or property damage or other type of loss resulting from the handling, storage or use of this product. The buyer assumes all risk and liability therefor and accepts and uses this product on these conditions.

Agrico Chemical Company
Memphis, Tennessee 38101

SUBSTITUTES:

Agrico Lawn Fungicide

BEST AVAILABLE COPY

2/2