

S A - 5 0
BRAND

40% Chlordane WETTABLE SPRAY CONCENTRATE

ACTIVE INGREDIENTS:
Technical Chlordane 40.0%
INERT INGREDIENTS 60.0%
100.0%

Equivalent to 14% Octachloro-1, 7-Methanotetrahydroindane and 16% related compounds.

KEEP OUT OF REACH OF CHILDREN

CAUTION: Harmful if swallowed. Contact with skin can cause toxic symptoms. Do not breathe dust or spray mist. Avoid contact with eyes, skin, and clothing. In case of contact with skin, wash with soap and water. For eyes, get medical attention. Avoid contamination of food and food stuffs. Do not reuse empty container. Destroy when empty.

NOTE: Dosages indicated are suggested only, and may not give satisfactory control under all conditions.

MIXING: Mix thoroughly with suggested amount of water. Agitate while spraying to hold in suspension.

Compatible with most insecticides and fungicides except oils and alkaline materials.

DO NOT feed plants treated with chlordane, or ensilage made from treated plants to poultry, dairy animals, or animals being finished for slaughter. Avoid spray drift to adjacent crops and pastures.

Highly toxic to fish and wildlife. Do not allow spray or permit drift to any body of water or apply to areas where crops not indicated on the label.

Buyer assumes all risks of the storage or handling of this material not in strict accordance with the directions given hereon.

Southern Agricultural Insecticides, Inc.
P.O. Box 100, Memphis, Tennessee 38101

SUGGESTED DOSAGES

NOTE: In certain areas, insects, particularly wireworms, have shown resistance to chlordane treatments. Do not apply wettable chlordane before obtaining from your state agricultural extension service additional information consistent with the label directions and caution.

FOR LAWN APPLICATIONS

Apply the suggested quantity of SA 50 Brand Chlordane Wettable Spray with sufficient water to obtain thorough coverage to each 1,000 square feet of lawn or turf area.

ANTS—Two level cupfull to entire area of termite mounds. Sprinkle after application.

ARMYWORMS—**FALL ARMYWORMS:** One level cupfull. Apply as soon as infestation appears—while worms are small.

BILLBUGS—Four level cupfull—when damage is observed.

JAPANESE BEETLE GRUBS—Two level cupfull—when examination of turf show grubs are present. Apply when ground is not frozen and wash into soil.

MOLE CRICKETS—One level cupfull—as soon as damage is noted.

SOD WEBWORMS—One level cupfull—in late afternoon or evening. A second application may be necessary.

CAUTION: Keep children and pets off treated lawns until insecticide has been washed off by rain or sprinkling, and grass is completely dry. Birds feeding over treated area may be harmed.

FOOD CROPS

MOLE CRICKET control in Broccoli, Cabbage, Cauliflower, Eggplant, Lettuce, Peppers, and Tomatoes.

SOIL BEDS—Drench using 5 oz. per 100 gal. of water per 1,000 sq. ft. of bed. **IN FIELD:** Before seeding or transplanting, apply to soil in enough water to obtain soil coverage using 3 to 5 lbs. per acre.

ROOT MAGGOTS in cabbage. Spray to the soil at the base of the plants, immediately following transplantation, using 3 to 5 lbs. per acre in enough water to obtain thorough coverage.

CORN ROOTWORMS control in corn. Apply 2 to 3 lbs. per acre before planting. Use 2 to 3 lb. rate for row treatment, 3 to 4 lb. rate for broadcast treatment. Immediately work into top 3 to 4 inches of soil.

WIREWORMS control in Broccoli, Cabbage, Cauliflower, Eggplant, Lettuce, Peppers, and Tomatoes. Broadcast prior to planting using 5 lbs. per acre. Thoroughly work into upper 6 to 9 inches of soil. One application should give control for 3 or more years.

CAUTION: Do not apply to food crops after edible parts have begun to form.

ORNAMENTALS

Apply as a full cover spray at rate of 1 level cupfull per gallon of water. Do not spray into open blooms.

ARBORVITAE AND JUNIPERS—to control bagworm. Apply in spring when insects first appear.

CAMELIA, HOLLY AND ELM—to control leaf beetle. Apply when insects or injury is noted.

ROSES AND PRIVET—to control thrips. Avoid spraying thrips on roses. Repeat as needed.

ROSES AND PRIVET—to control thrips. Avoid spraying thrips on roses. Repeat as needed.

