

PM 23 14

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

NOV 29 1989

Mr. Ronald L. Costa, Inc.  
Southern Agricultural Industries, Inc.  
P.O. Box 211  
Lafayette, La. 70508

Dear Mr. Costa: Re:

Subject: [Faint text regarding agricultural products and EPA regulations]

A stamped copy is enclosed for your records.

**BEST AVAILABLE COPY**

55491; I:Vohra:MF-12; KENCO: 11/27/89; 09/90; DD:vo;jh:dg

CONCURRENCES							
SYMBOL	H7505C						
SURNAME	S. VOHRA						
DATE	11/28/89						

4. Please submit three (3) copies of your final printed labeling before you release the product for shipment.

Sincerely yours,

Joanne I. Miller  
Acting Product Manager (23)  
Fungicide-Herbicide Branch  
Registration Division (H7505C)

Enclosure

BEST AVAILABLE COPY

GREEN BORDER  
YELLOW BACKGROUND

SA 50 TOMATO DUST

ACTIVE INGREDIENT:	By. Wt.
BASIC COPPER SULFATE* .....	13.30%
INERT INGREDIENTS:.....	86.70%
TOTAL:.....	100.00%
*Metallic Copper Equivalent.....	7.19%

KEEP OUT OF REACH OF CHILDREN 12 FT

CAUTION 18 FT

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush eyes with plenty of water. Seek medical attention as soon as possible if irritation persists. IF INHALED: Remove victim to fresh air. Contact a physician if irritation occurs. IF SWALLOWED: Give victim 1 or 2 glasses of water and induce vomiting by touching back of throat with finger or blunt object. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention. IF ON SKIN: Wash affected area with plenty of soap and water. Contact a physician if irritation occurs.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS. CAUTION: HARMFUL IF SWALLOWED. AVOID CONTACT WITH EYES, SKIN, OR CLOTHING. AVOID BREATHING DUST. WASH THOROUGHLY WITH SOAP AND WATER AFTER HANDLING. MAY CAUSE SKIN SENSITIZATION REACTIONS IN CERTAIN INDIVIDUALS.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic organisms. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not contaminate water by cleaning of equipment or disposal of wasters.

EPA Est. No. 829-F1-1(3): 829-NC-1(4): 829-NC-2(5) Number following Est. No. corresponds to first digit in Lot No. on bottom of bag.

SOUTHERN AGRICULTURAL INSECTICIDES, INC.

Palmetto, FL\* 34220  
EPA Reg. No. 829-6

Hendersonville, NC 28793  
J2

Boone, NC 28607  
Net Weight: 5 pounds

BEST AVAILABLE COPY

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

NOV 29 1989

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, this pesticide is registered under FIFRA No. 829-6

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**A FUNGICIDE DUST FOR USE ON:**

**TOMATOES and POTATOES** for the control of early and late blights and as an aid in controlling bacterial spot. Dust plants thoroughly at weekly intervals and following rains. For potatoes make 30 to 60 lbs. per acre applications beginning when plants are 6 inches high. For tomatoes make 30 to 60 lbs. per acre applications beginning when plants are set in the field.

**CUCUMBERS, SQUASH, WATERMELONS and CANTALOUPES** for the control of downy mildew. Dust thoroughly, but lightly, using 15 to 40 lbs. per acre. Apply at 10 day intervals and following rains. Heavy applications, applications to tender foliage, and applications to wet foliage can cause injury to cucurbits.

**PEPPER** as an aid in the control of bacterial spot, apply at weekly intervals and if possible preceding heavy rains, using 20 to 45 lbs. per acre applications. There is no time limitation on the use of this dust. Remove excess residue at harvest. Consult proper State Extension specialist to determine if local recommendations are consistent with this label before using. Do not exceed indicated dosages. May cause injury to tender plant parts particularly when applications are excessive, uneven, or when foliage is wet. Control of the indicated diseases may not be satisfactory during periods of frequent or heavy rainfall, when disease infestations are extremely severe, or when applications are uneven, inadequate, or poorly timed.

Do not apply or allow drift to corn, grain sorghum, or other copper sensitive plants. Do not apply when conditions favor drift from treated area.

**STORAGE AND DISPOSAL 12 PT**

**PESTICIDE STORAGE:** Store pesticide in the original container in a locked storage area out of the reach of children and pets.

**PESTICIDE DISPOSAL:** To dispose of pesticide, securely wrap original container in several layers of newspaper and discard in trash.

**CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

Buyer assumes all risks of use, storage, or handling of this material in strict accordance with the directions given herewith.

