

# SA BRAND 50<sup>®</sup> Ferbam Dust

A protective fungicide for the control of

## Blue Mold in tobacco plant beds

**ACTIVE INGREDIENTS:**

Ferbam (ferric dimethyl dithiocarbamate) ..... 11.40%

**INERT INGREDIENTS** ..... 88.60%

**100.00%**

Begin applications when plants are about the size of a dime. Apply at 3-day intervals, until plants are set in the field. Make 2 to 3½ lbs. (per 100 square yards of bed) applications depending on the size and spacing of the plants. Repeat applications following rains. When Blue Mold is detected in the bed increase application frequency to 3 times per week.

Thoroughness of application is essential for Blue Mold control. Plants should be inspected after dusting to confirm thoroughness of coverage. May cause injury to tender plants especially when application is uneven or when foliage is damp. Control of Blue Mold may not be satisfactory during periods of heavy rainfall, when disease infestations are extremely severe, or when applications are uneven, inadequate, or poorly timed.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply when conditions favor drift from areas treated. Apply this product only as specified on this label. Do not contaminate water by cleaning of equipment or disposal of wastes.

### Keep Out of Reach of Children

**CAUTION!** May cause irritation of eyes, nose, throat and skin. Avoid breathing dust. Avoid contact with eyes, skin or clothing. In case of contact, flush with plenty of water; for eyes get medical attention. Keep dust dry. Store in a safe dry place. Keep away from fire and sparks.

Buyer assumes all risks of use, storage, or handling not in strict accordance with the precautions and directions given herewith.

## Southern Agricultural Insecticides, Inc.

Palmetto, Florida      Boone, N. C.      Hendersonville, N. C.

U. S. D. A. Reg. No. 829-28      C

NET WEIGHT

LABEL WILL BE TRIMMED TO 4" X 6"

