SA BRAND 50¢

5% Malathion Dust

ACTIVE INGREDIENTS:
Malathion (O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate) 5.00%
INERT INGREDIENTS 95.00%

100.00%

Keep Out of Reach of Children

CAUTION! Harmful if swallowed. Do not breathe dust. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling. Do not contaminate feed or foodstuffs. Burn empty container. Stay out of smoke and fumes. Bury waste dust in an isolated area.

Highly toxic to fish. Do not contaminate any body of water. Do not apply when conditions favor drift from treated area.

NOTE: Apply only to dry foliage. Dosages given are suggested only and may not give satisfactory control under all conditions. Where dosage ranges are indicated, apply larger dosages to maturing crops with compact spacing, proportionately less for smaller plants and wider spacing. Do not exceed suggested dosages.

BEANS for the control of aphids, Mexican bean beetle and leafhoppers. Apply 25 to 35 lbs. per acre as applications as needed. Do not apply within 24 hours of harvest.

BROCCOLI, CABBAGE, CAULIFLOWER, COLLARDS, KALE, MUSTARD and TURNIPS (in home garden) for the control of aphids and imported cabbage worm. Apply 25 to 50 lbs. per acre every 7 days beginning when first true leaves appear. Do not apply to broccoli or turnips within 3 days of harvest. Do not apply to cabbage, cauliflower, collards, kale or mustard within 7 days of harvest.

CUCUMBERS and SQUASH (in home gardens) to control aphids and cucumber beetle adults. Apply [illegible] lbs. per acre to foliage as needed. Do not apply to cucumbers and squash within 1 day of harvest.

PEAS: GARDEN, BLACK EYED and COWPEAS for the control of aphids, apply 25 lbs. per acre in applications as needed. Do not apply within 3 days of harvest. If vines are to be used for animal feed do not apply within 7 days of harvest.

(SEE OTHER PANEL FOR FURTHER INSTRUCTIONS)

Southern Agricultural Insecticides, Inc.

PALMETTO, FLA. BOONE, N. C. HENDERSONVILLE, N. C.

NET WT. 5 LBS.

PEPPER (in home garden) for the control of aphids, apply 20 to 30 lbs. per acre to foliage as needed. Do not apply to peppers within 3 days of harvest.

TOMATOES to control aphids and spider mites, apply 25 to 40 lbs. per acre to foliage as needed. Do not use on tomatoes within 1 day of harvest.

STRAWBERRIES to control aphids, [illegible] potato [illegible] spider mite, and cattle lice, apply 25 to 50 lbs. per [illegible] as needed. Do not apply to ripe berries within 3 days [illegible] harvest. Do not use Malathion where cyclamen mite is a problem.

TOBACCO for aphids in plant beds apply 3/4 to 1 lb. per 100 square yards of bed. Apply with thorough, even dusting, [illegible] as necessary. Do not apply within 3 days of transplanting.

Apply to the following Ornamentals using different dust to obtain light but thorough, even coverage.

ASTERS to control aphids; apply as needed.

AZALEA to control azalea lace bug and spider mite, make 3 applications 7 days apart.

GARDENIA to control white fly; repeat at 10-day intervals as necessary.

PYRACANTHA to control hawthorne lace bugs; apply as needed.

ROSES for the control of aphids, leafhoppers and thrips; apply at 3-day intervals as needed.

VIOLETS to control mealy bugs; apply as needed.

CAUTION: Applications to ornamentals in bloom may cause injury to blooms. Do not apply to Hibiscus, Papaya, certain ferns including Boston, Maiden Hair, and Pteris, as well as some species of Crassula.

HOME GARDEN CITRUS: Oranges, Grapefruit, Lemons, Limes, Tangelos, and Tangerines for the control of purple scale, red scale, soft brown scale and aphids, apply with thorough light dusting. Repeat as necessary. Do not apply while plants are in bloom. Do not apply within 7 days of harvest.

CAUTION: Do not apply to very young or weak animals. Do not apply in conjunction with oral drenches, internal medications, or other organic phosphate treatments. Do not contaminate feed, water, feed and watering utensils, milk, or milking equipment.

DOGS for the control of fleas, dust animals thoroughly with a puff type duster. If dog is confined, one application may suffice. If animal is allowed to run free reinfestation may occur. Application may be repeated in 3 weeks. Do not use on puppies or dogs that are in weakened condition. To prevent reinfestation from such areas as pens, kennels and under porches, remove manure and debris, and dust at rate of 2 lbs. per 1,000 square feet, using hand duster; repeat as necessary.

BEEF CATTLE to control lice, treat animals thoroughly and repeat after 2 or 3 weeks if needed. Do not treat [illegible] dairy animal in this manner.

DAIRY AND BEEF CATTLE to control lice, apply 1 oz. per animal to the back, neck and thighs. Repeat applications at 10 to 14 day intervals as needed.

Do not apply to dairy cattle during milking or within 3 [illegible] of milking.

HOGS to control lice, treat animals thoroughly, repeat in 2 or 3 weeks if needed.

POULTRY for the control of Northern Fowl Mite and Lice. Apply 1 lb. to 100 square feet of litter.

[illegible] handling [illegible] with [illegible]