

DEC 19 1972

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 829-67 SUBJECT
TO ATTACHED COMMENTS.

SA BRAND 50[®] MANEB DUST

ACTIVE INGREDIENTS:

Maneb (manganese ethylenebisdithiocarbamate) 5.6%
(Manganese as equivalent metallic 1.16%)

INERT INGREDIENTS 94.4%

— A FUNGICIDE DUST FOR USE ON — 100.0%

Beans as an aid in the control of red rust. Apply when disease threatens, using 25 to 50 lbs. per acre applications depending on size of plants. Repeat applications at 5 to 7-day intervals. Do not apply within 4 days of harvest.

Cabbage for the control of Alternaria leaf spot and downy mildew. Plant beds: make 40 lbs. per acre (1 lb. per 1000 sq. ft. of bed) applications at 3-day intervals, until plants are set in field. Field application: make 40 lbs. per acre applications at 5 to 7-day intervals when disease threatens. Do not apply to cabbage within 7 days of harvest.

Cucumbers, squash, and melons for the control of downy mildew, anthracnose and scab. Begin application when true leaves appear. Make 25 to 50 lbs. per acre applications at weekly intervals and following rains. For scab control in the Southern Appalachian region apply twice weekly when disease threatens. Do not use within 5 days of harvest.

Tomatoes and potatoes for the control of gray leaf spot, early and late blights ^{blight}. Begin applications when tomatoes are set in field and when potatoes are 2 to 6 inches in height. Apply at weekly intervals and following rains using 30 to 40 lbs. per acre applications. For control of late blight on maturing tomatoes and potatoes make 50 lbs. per acre applications when applied by ground duster, and 60 lbs. when applied by aircraft. On tomatoes, do not use within 5 days of harvest. Remove excess residue of dust at harvest.

Gladiolus for the control of Botrytis, apply 30 to 40 lbs. per acre applications at 5 to 7-day intervals beginning when disease threatens. May cause injury to tender plant parts especially when application is uneven or when foliage is damp. Control of the indicated diseases may not be satisfactory during periods of frequent and heavy rainfall, when disease or insect infestations are extremely severe or when applications are uneven, inadequate or poorly timed.

Keep Out of Reach of Children

CAUTION! May be harmful if swallowed. May irritate eyes, nose, throat, and skin. Avoid breathing dust. Avoid contact with skin and eyes. Wash thoroughly after handling. Do not contaminate feed or foodstuffs. Store in a safe dry place. Burn empty container. Stay out of smoke and fumes. Toxic to fish. Do not contaminate any body of water nor allow drift from the crop area treated.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with the directions and precautions given herewith.

Southern Agricultural Insecticides, Inc.

PALMETTO, FLA. BOONE, N. C. HENDERSONVILLE, N. C.
EPA Reg. No. 829-67 C