

S A - 5 0° BRAND 10% CHLORDANE GRANULES

ACTIVE INGREDIENTS:
Technical Chlordane (including 6.0% octachloro-
4,7-methanotetrahydroindane and 4.0% related compounds) 10.0%
INERT INGREDIENTS 90.0%

KEEP OUT OF REACH OF CHILDREN 100.0%

CAUTION: Harmful if swallowed. Contact can cause toxic symptoms. Avoid breathing dust. In case of contact with skin wash with soap and water. Avoid contamination of feed and foodstuffs.

ANTODOTE: External - In case of spillage on skin, wash promptly with soap and water. Internal - Emetic 1 tablespoon of mustard in tumbler of water. Call a physician.

CONTAINER DISPOSAL: Do not reuse empty container. Rinse equipment and containers and dispose of wastes by burying in noncrop-lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes or by burning. Keep out of smoke.

This product is toxic to fish, birds, and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

For use on noncropland and non-food crop area: such as sod, ornamental turf, industrial sites, roadsides and ornamental plant nurseries in areas covered by Federal or State quarantine programs for the two insects listed below. These treatments may be used to certify to pest free conditions as well as for use in suppression and control programs.

FIRE ANTS: BROADCAST SOIL TREATMENT - Apply this material uniformly over surface using 10 to 15 lbs. per acre (1/4 to 1/3 lb. per 1,000 sq. ft.).

MOUND APPLICATION: Apply 3/4 to 1 1/2 lbs of this product to the entrance of the nest as well as to an area 3 feet around the mound. Treated areas should be examined, 3 weeks after treatment and surviving colonies retreated. Do not apply when soil is wet.

JAPANESE BEETLE LARVAE: Apply this material uniformly to surface at the rate of 50 - 100 lbs. per acre (1 1/8 - 2 1/4 lbs. per 1,000 sq. ft.). Apply just before rain or use sprinklers to remove material from turf.

DO NOT APPLY when people or animals are on lawn. Keep children and pets off treated areas until Chlordane has been washed off grass by sprinkling and grass has completely dried.

DO NOT APPLY to pastures or range land or graze livestock on treated areas within 1 year of application.

THIS MATERIAL may not give satisfactory control under all conditions. Timing and method of application, weather and growing conditions and other influencing factors in the use of this material are beyond the control of the seller. Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with the directions and precautions given herewith.

EPA EST 829-F 1-1 (3); 829-NC-1 (4); 829-NC-2 (5) Number following Est. No. Corresponds to first digit in Lot No. on bottom of bag.

Southern Agricultural Insecticides, Inc.

Palmetto, Fla. Boone, N.C. Hendersonville, N.C.

EPA Reg. No. 829-77 E Net Weight 1 lb.

