

S
SO

Do not apply from lack of not apply at sulfur or in temperature

ACTIVE INGREDIENTS
Petroleum (Unsulfurated)
INERT INGREDIENTS

The oil in this

CAUTION

In general, tender succulent plants, they applying oil and continue stand.

For scale control insects are Do not use on plant injury conditions. Follow instructions in the use assumes all with the directions.

South

Polmetto, Fla.
EPA Reg. No.

NON-DORMANT USES

SUGGESTIONS FOR USE WHEN PLANT IS ACTIVELY GROWING

This includes all suggested Florida uses as well as suggested uses on actively growing plants (non-dormant) in other Southeastern areas. Do not apply in Florida after October 1st. Do not use when temperatures fall below 42 F. within 3 weeks of application. If second application of oil is necessary, do not apply within six weeks after first oil spray. Avoid spraying open blooms.

829-83

PLANT	INSECT	Quarts per 100 gal.	Tablespoonfuls per gal.	
Avocado	Wax Scale	5	3	Preferably in May, June, or July.
	White Fly			
Aloe	Latania Scale	5	3	2 applications after blooming period.
	Aloe Lace Bug			
	White Fly Larva			
Camellia	Scales	5	3	2 applications at close of blooming period.
	White-Fly Larva			
	Citrus Mealy Bug			
Carni	Soft Scales	5	3	Preferably in June or July.
	Florida Red Scale			
	Spider Mite			
Citrus	White Fly Larva	4	2	2 applications as needed.
	Soft Scales			
Clematis	Mealy Bug	5	3	2 applications, preferably after blooming period.
	White Fly Larva			
Holly	Soft Scales	5	3	2 applications.
	White Fly Larva			
Hibiscus	White Fly Larva	4	2	2nd application only if needed.
Ixora	Soft Scales	5	4	2 applications as needed, preferably after blooming.
	Mealy Bug			
Ivy	Oleander Scale	2	1	Avoid 2nd application. Preferably spring or fall.
Jasmine	White Fly Larva	5	3	2 applications as needed.
	Soft Scales			
Mango	Florida Red Scale	5	3	2nd application only if needed.
	Mango Scale			
Palm, Coccothraupis	Soft Scales	5	2	Rinse with clear water 1 to 2 hours after applying.
	Mealy Bug			
Pittosporum	Cottony Cushion Scale	5	3	2 applications as needed.
Viburnum	Soft Scales	5	3	2 applications as needed.

DORMANT USES

Dormant oil sprays are applied during the winter and early spring in those areas where many plants reach true dormancy (not Peninsular Florida).

Dormant oil sprays should not be applied in freezing weather. Preferably, temperatures should be above 40 F.

Dormant sprays are generally more effective in the spring just before new growth starts.

Apple	Scale	5	3	Before visible growth starts.
Apple	Black Scale	5	3	On early blooming varieties, apply before starting warm weather in late fall.
Camellia	Scales	5	3	Same as for active growth.
Holly	Holly Scale	8	5	1 application before new growth starts.
Pittosporum	Cottony Cushion Scale	5	3	2 applications.
Pine	White Pine Scale	12	7	1 application 2 weeks after new growth starts.
Pine	Wax Scale	12	7	Before visible growth starts.
Pine	Pine Scale	6	4	Before visible growth starts.

AS A PLANT SHINE

For a bright, clear, lustrous shine on plants. Spray or wipe with a soft cloth or sponge with 2 to 3 tablespoonfuls per gallon of water.