

829-83

05-16-2012

1
10



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

MAY 16 2012

Dr. Matthew Brooks, Director
Ag-Chem Consulting
Authorized Representative for Southern Ag. Inc
12208 Quinque Lane
Clifton, VA 20124

S916968
300/1155

Dear Dr. Brooks:

Subject: SOLUBLE OIL SPRAY
EPA Registration Number 829-83
Label Submission dated May 15, 2012

The proposed amended labeling coded "5-3-12" is accepted under the provisions of the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended.

When labels are printed, the following changes should be made.

Commercial Use - Sublabel A

1. In the second sentence of the "EFFICACY CONSIDERATIONS" paragraph (page 3), "make" should be changed to "made".

Residential Use - Sublabel B

1. In the "Note" entry for "Palm" in the table below the "NON-DORMANT USES WHEN PLANT IS ACTIVELY GROWING" paragraph (page 7), "Rinse with clear water 1 ½ to 2" should be changed back to "Rinse with clear water 1 ½ to 2 hours after applying". The "Palm" designation itself should be changed back to "Palm, outdoor".
4. The registrant's name and address(es), the product's registration number, and the production establishment's number must appear on "**Sublabel B**" as it is to be a completely separate label from "**Sublabel A**". Both "sublabels" also should include the registrant's telephone number.

On page 8 of the label accepted via this letter, there is a reference to two web addresses linked to this product's registrant. Consequently, those websites become labeling under the Federal Insecticide Fungicide and Rodenticide Act and are subject to review by the Agency. If either website included statements or other representations that are false or misleading in any particular, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) provides examples of statements EPA considers to be false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, if the Agency finds or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.


A stamped copy of the label is enclosed for your records.

Submit two copies of final printed labeling before releasing this product for shipment under the labeling accepted via this letter.

If these conditions are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6(e).

If you have any questions regarding this letter, contact Dr. William W. Jacobs by phone at 703-305-6406, by FAX at 703-308-0029, or via e-mail at jacobs.bill@epa.gov.

Sincerely yours,



Venus Eagle
Product Manager 01
Insecticide-Rodenticide Branch
Registration Division (7505P)

3
10

Intended for Commercial Use-Sublabel A

SOLUBLE OIL SPRAY

SUPERIOR 70-SECOND TYPE

FOR USE ON CITRUS TREES, APPLE TREES, PALM TREES, WOODY ORNAMENTALS (as listed on this label), AND VEGETABLES (as listed on this label) TO CONTROL WHITE FLIES, SCALE INSECTS, MEALY BUGS, AND OTHER TYPES OF INSECTS, AS LISTED ON THIS LABEL.

See DIRECTIONS FOR USE

ACTIVE INGREDIENT

Mineral Oil 98.00%
(Unsulfonated residue not less than 92%)

INERT INGREDIENTS: 2.00%

TOTAL: 100.00%

Contains petroleum distillates

The oil in this product meets Florida Citrus Standards FC 435.66

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

ACCEPTED
MAY 16 2012

Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended, for the
pesticide registered under:

EPA. Reg. No: 829-83

received 5/16/12

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give any liquid to the person. • Do not give anything by mouth to an unconscious person
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for treatment advice.
<p>Have product container or label with you when calling a poison control center or doctor or going for treatment. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378</p>	
<p>Note to Physician: Contains petroleum distillate and may pose an aspiration pneumonia hazard.</p>	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled, or absorbed through the skin. Do not get in eyes, on skin, or on clothing. Avoid breathing spray mist. Wash contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category E on an EPA chemical resistance category selection chart. Applicators and other handlers must wear:

1. Long-sleeved shirt and long pants.
2. Chemical resistant gloves, made from barrier laminate, nitrile rubber, neoprene rubber, or viton.
3. Shoes plus socks.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent material that have been drenched or heavily contaminated with the products concentrate. Do not reuse them.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

5
10

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Note: Do not apply this product through any type of Irrigation System.

Aerial application of this product is prohibited. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this labeling about personal protective equipment (PPE), and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- (1) Coveralls,
- (2) Chemical-resistant gloves, made of any waterproof material, and
- (3) Shoes plus Socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Do not enter or allow others to until spray has dried.

RESTRICTIONS AND PRECAUTIONS

Do not apply this product in a way that will contact workers or others persons, either directly or through drift. Only protected handlers may be in the area during application. Mix thoroughly the proper amount of oil and water in the spray tank and continue agitation during spraying. In general, woody ornamentals are more tolerant to oil sprays than are tender succulent plants. If oil applications must be made to tender plants, they may be syringed off with clear water about an hour after applying oil.

Do not apply oil sprays to plants that are wilting or are near wilting from lack of moisture. Do not apply oil with sulfur or lime sulfur. Do not apply an oil spray within 3 weeks before or after an application of sulfur or lime sulfur. Avoid spraying during periods of excessively high temperature (above 90°F). Do not exceed three oil sprayings per year. Do not use more than the specified dosage, for to do so may cause plant injury. Do not allow the mixture to stand.

EFFICACY CONSIDERATIONS

For scalcicides to be effective, thorough coverage is essential. Better scale control can be anticipated when applications are made when scale insects are in the crawler stage.

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g. ground, airblast) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Wind Speed: Do not apply at wind speeds greater than 15 mph at the application site.

Temperature Inversions: If applying at wind speeds less than 3 mph, the applicator must determine if a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions

Droplet Size: Apply as a medium or coarser spray (ASABE standard 572), and the minimum volume mean diameter (VMD) for spinning atomizer nozzles.

Ground Applications: Apply using a nozzle height of no more than 4 feet above the ground or canopy. Applications must be consistent with the limitations for wind speed, temperature inversions, and droplet size indicated above.

Airblast Applications: For airblast applications, turn off outward-pointing nozzles at row ends and when spraying outer row. To minimize spray loss over the top in orchard applications, spray must be directed into the canopy. Applications must be consistent with the limitations for wind speed, temperature inversions, and droplet size indicated above.

NON-DORMANT USES (WHEN PLANT IS ACTIVELY GROWING)

This includes all Florida uses as well as uses on actively growing plants (non-dormant) in other Southeastern areas. Do not use when temperatures fall below 42°F within 3 weeks of application. If a second application of oil is necessary, do not apply within six weeks after first oil spray. Avoid spraying open blooms.

PLANT	INSECT	Qts. per 100 Gal.	Tbls. per Gallon	
Azaleas	Azalea Lace Bug, White Fly Larva	5	3 1/2	2 applications after blooming period.
Camellia	Scales, White Fly Larva, Citrus Mealy Bug, Soft Scales	5	3 1/2	2 applications at close of blooming period.
Citrus	Florida Red Scale, Purple Scale, Spider Mite, White Fly Larvae.	5	4	Preferably in June or July.
Crotons	Soft Scales, Mealy Bug	4	2	2 applications, if needed.
Gardenia	White Fly Larvae, Soft Scales, Citrus Mealy Bug	5	3 1/2	2 applications preferably after blooming period.
Holly	White Fly Larvae	5	3 1/2	2 applications.
Hibiscus	White Fly Larvae	4	2	Make a second application only if needed.
Ixora	Soft Scales, Mealy Bug	5	4	2 applications as needed, preferably after blooming.
Ivy	Oleander Scale	2	1	Make only one application per year, preferably spring or fall.
Ligustrum	White Fly Larva Soft Scales	5	3 1/2	Make a second application only if needed.
Mango	Florida Red Scale, Mango Scale	5	3 1/2	Make a second application only if needed.
Palm, outdoor	Soft Scales, Mealy Bug	3	2	Rinse with clear water 1 1/2 to 2 hours after applying.
Pittosporum	Cottony Cushion Scale	5	3	2 applications as needed.
Viburnum	Soft Scale	5	3	2 applications as needed.

For application to Citrus, do not exceed 159 pounds of active ingredients per acre, in a maximum of 1,500 gallons of spray mix.

For use on the types of vegetables listed below, do not exceed 3 applications in a growing season and do not apply at intervals of less than 14 days. On these vegetable crops, this product controls Aphids, Mites, Beetle Larvae, Leafminers, Thrips, Leafhopper, and Whitefly (except for avocado as noted in table below).

PLANT	INSECT	Qts. per 100 Gal.	Tbls. per Gallon
Avocado	Wax Scale, White Fly, Latania Scale	5	3 1/2
Beans		4-8	2 1/2 - 5
Beets		4-8	2 1/2 - 5
Cabbage		4-8	2 1/2 - 5
Cauliflower		4-8	2 1/2 - 5
Celery		4-8	2 1/2 - 5
Corn		4-8	2 1/2 - 5
Cucurbit		4-8	2 1/2 - 5
Eggplant		4-8	2 1/2 - 5
Melon		4-8	2 1/2 - 5
Pepper		4-8	2 1/2 - 5
Squash		4-8	2 1/2 - 5
Tomato		4-8	2 1/2 - 5
Tobacco		4-8	2 1/2 - 5

DORMANT USES

Dormant oil sprays are applied during the winter and early spring in those areas where many plants reach true dormancy (not Peninsular Florida). Dormant oil sprays should not be applied in freezing weather. Preferably, temperatures should be above 40°F. Dormant sprays are generally more effective in the spring just before new growth starts.

PLANT	INSECT	Qts. per 100 Gal.	Tbls. per Gallon	
Apples	Scales	8	6	Before visible growth starts.
Azalea	Bark Scales	6	4	On early blooming varieties, apply during warm weather in late fall.
Camellia	Scales	6	4	Supplement to summer applications.
Holly	Holly Scale	8	6	In spring before buds open.
Pittosporum	Cottony Cushion Scale	8	6	2 applications.
Privet	White Peach Scale	12	8	2 applications 2 weeks apart.
Pyracantha	Wax Scale	12	8	Before visible growth starts.
Rose	Rose Scale	6	4	Before visible growth starts

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

(FOR USE ON CONTAINERS 5 GALLONS OR LESS)

PESTICIDE STORAGE: Open dumping is prohibited. Store only in original container. Do not reuse empty container. If a leaky container must be contained within another, mark the outer container to identify the contents. Store pesticides away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Keep this product under locked storage sufficient to make it inaccessible to children or persons unfamiliar with its proper use.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incinerate if allowed by State and local authorities.

(FOR USE ON REFILLABLE 30, 55, 275 GALLON CONTAINERS)

PESTICIDE STORAGE: Open dumping is prohibited. Store only in original container. Do not reuse empty container. If a leaky container must be contained within another, mark the outer container to identify the contents. Store pesticides away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Keep this product under locked storage sufficient to make it inaccessible to children or persons unfamiliar with its proper use.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Refillable container. Refill this container with this product only. Do not use this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Offer container for recycling or reconditioning, if available, or dispose of empty container in a sanitary landfill or by incineration, if allowed by State and local authorities.

WARRANTY STATEMENT

Follow instructions carefully. Timing, method of application, weather conditions, mixtures with other chemicals and other influencing factors in the use of this product are beyond control of the seller.

To the extent consistent with applicable laws buyer assumes all risks of use, storage, or handling not in strict accordance with the directions and precautions given herewith.

Southern Agricultural Insecticides, Inc.

PALMETTO, FL. 34220

HENDERSONVILLE, N.C. 28793

BOONE, N.C. 28607

Phone: 941-729-4534

EPA Reg. No. 829-83

EPA Est. No. 829-FL-1

Net Contents Liquid:

1 pint	1 Gallon	2.5 Gallons	30 Gallons	50 Gallons	275 Gallons
(0.473 liters)	(3.785 liters)	(9.46 liters)	(113.55 liters)	(189 liters)	(1041 liters)

8/10

SOLUBLE OIL SPRAY
Intended for Residential Use –Sublabel B

FOR USE ON CITRUS TREES, APPLE TREES, PALM TREES, WOODY ORNAMENTALS (as listed on this label), AND VEGETABLES (as listed on this label) TO CONTROL WHITE FLIES, SCALE INSECTS, MEALY BUGS, AND OTHER TYPES OF INSECTS, AS LISTED ON THIS LABEL.
See DIRECTIONS FOR USE

FOR RESIDENTIAL USE ONLY

ACTIVE INGREDIENT

Mineral oil* 98.00%
(Unulfonated residue not less than 92%)

INERT INGREDIENTS: 2.00%

TOTAL: 100.00%

*Contains petroleum distillates

The oil in this product meets Florida citrus Standards FC 435.66

KEEP OUT OF REACH OF CHILDREN
CAUTION

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give any liquid to the person. • Do not give anything by mouth to an unconscious person
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for treatment advice.
<p>Have product container or label with you when calling a poison control center or doctor or going for treatment. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378</p>	
<p>Note to Physician: Contains petroleum distillate and may pose an aspiration pneumonia hazard.</p>	

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled, or absorbed through the skin. Do not get in eyes, on skin, or on clothing. Avoid breathing spray mist. Wash contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Wear a long-sleeved shirt, long pants, socks and waterproof gloves while using or handling this product.

5/12

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Note: Do not apply this product through any type of Irrigation System.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

USE RESTRICTIONS AND PRECAUTIONS

Mix thoroughly the proper amount of oil and water in the spray tank and continue agitation during spraying. Do not allow the mixture to stand. In general, woody ornamentals are more tolerant to oil sprays than are tender succulent plants. If oil applications must be made to tender plants, they may be syringed off with clear water about an hour after applying oil.

Do not apply oil sprays to plants that are wilting or are near wilting from lack of moisture. Do not apply oil with sulfur or lime sulfur. Do not apply an oil spray within 3 weeks before or after an application of sulfur or lime sulfur. Avoid spraying during periods of excessively high temperature (above 90°F). Do not exceed three oil sprayings per year. Do not use more than the specified dosage, for to do so may cause plant injury.

HOW TO IMPROVE CONTROL

For scalicides to be effective, thorough coverage is essential. Better scale control can be anticipated when applications are made when scale insects are in the crawler stage.

NON-DORMANT USES (WHEN PLANT IS ACTIVELY GROWING)

This includes all Florida uses as well as uses on actively growing plants (non-dormant) in other Southeastern areas. Do not use when temperatures fall below 42°F within 3 weeks of application. If second application of oil is necessary, do not apply within six weeks after first oil spray. Avoid spraying open blooms.

PLANT	INSECT	Tbls. per Gallon	Note
Azaleas	Azalea Lace Bug, White Fly Larva	3 1/2	2 applications after blooming period.
Camellia	Scales, White Fly Larva, Citrus Mealy Bug, Soft Scales	3 1/2	2 applications at close of blooming period.
Citrus	Florida Red Scale, Purple Scale, Spider Mite, White Fly Larvae.	4	Preferably in June or July.
Crotons	Soft Scales, Mealy Bug	2	2 applications, if needed.
Gardenia	White Fly Larvae, Soft Scales, Citrus Mealy Bug	3 1/2	2 applications preferably after blooming period.
Holly	White Fly Larvae	3 1/2	2 applications.
Hibiscus	White Fly Larvae	2	Make a second application only if needed
Ixora	Soft Scales, Mealy Bug	4	2 applications as needed, preferably after blooming.
Ivy	Oleander Scale	1	Make only one application, Preferably spring or fall.
Ligustrum	White Fly Larva Soft Scales	3 1/2	Make a second application only if needed
Mango	Florida Red Scale, Mango Scale	3 1/2	Make a second application only if needed
Palm,	Soft Scales, Mealy Bug	2	Rinse with clear water 1 1/2 to 2
Viburnum	Soft Scale	3	2 applications as needed.

10
10

For use on the types of vegetables listed below, do not exceed 3 applications in a growing season and do not apply at intervals of less than 14 days . On these vegetable crops, this product controls Aphids, Mites, Beetle Larvae, Leafminers, Thrips, Leafhopper, and Whitefly (except for avocado as noted in table below).

PLANT	INSECT	Tbls. per Gallon
Avocado	Wax Scale, White Fly, Latania Scale	3 1/2
Beans		2 1/2 - 5
Beets		2 1/2 - 5
Cabbage		2 1/2 - 5
Cauliflower		2 1/2 - 5
Celery		2 1/2 - 5
Corn		2 1/2 - 5
Cucurbit		2 1/2 - 5
Eggplant		2 1/2 - 5
Melon		2 1/2 - 5
Pepper		2 1/2 - 5
Squash		2 1/2 - 5
Tomato		2 1/2 - 5
Tobacco		2 1/2 - 5

DORMANT USES

Dormant oil sprays are applied during the winter and early spring in those areas where many plants reach true dormancy (not Peninsular Florida). Dormant oil sprays should not be applied in freezing weather. Preferably, temperatures should be above 40°F. Dormant sprays are generally more effective in the spring just before new growth starts.

PLANT	INSECT	Tbls. per Gallon	Note
Apples	Scales	6	Before visible growth starts.
Azalea	Bark Scales	4	On early blooming varieties, apply during warm weather in late fall.
Camellia	Scales	4	Supplement to summer applications.
Holly	Holly Scale	6	In spring before buds open.
Pittosporum	Cottony Cushion Scale	6	2 applications.
Privet	White Peach Scale	8	2 applications 2 weeks apart.
Pyracantha	Wax Scale	8	Before visible growth starts.
Rose	Rose Scale	4	Before visible growth starts.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store pesticide in the original container in a locked storage area.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container.

If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

If empty: Place in trash or offer for recycling if available.

WARRANTY STATEMENT

Follow instructions carefully. Timing, method of application, weather conditions, mixtures with other chemicals and other influencing factors in the use of this product are beyond control of the seller.

To the extent consistent with applicable law buyer assumes all risks of use, storage, or handling not in strict accordance with the directions and precautions given herewith.

For information or questions contact our website at Southernag.com or email us at info@southernag.com

Southern Agricultural Insecticides, Inc.

PALMETTO, FL. 34220

HENDERSONVILLE, N.C. 28793

BOONE, N.C. 28607

Phone: 941-729-4534

EPA Reg. No. 829-83

EPA Est. No. 829-FL-1

Net Contents Liquid: 1 pint and 1 gallon