

Do not get in eyes, on skin, or on clothing. Do not breathe vapor or spray mist. Wash thoroughly with soap and water after handling and before eating or smoking. Wear clean clothing daily, clean synthetic rubber gloves and a mask or respirator of a type passed by the United States Bureau of Mines for Endrin protection. Shower and wash thoroughly and change to clean clothing after each day's spraying. Do not apply or allow drift to areas occupied by unprotected humans or beneficial animals. Do not contaminate feed or foodstuffs.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Not for use or storage in or around the home.

MAKE POSTHARVEST APPLICATION ONLY. Make single application to the orchard floor in October or November. Specific directions concerning nozzles should be obtained from State Agricultural Extension Specialist. Do not disturb trail system. Allow at least 2 months before and after treatment before cultivating or mowing.

PINE MOUSE (*pitymys pinetorum*) control. Spray must be applied at 500 lbs. per square inch pressure in order to penetrate ground cover to reach trail systems. Dilute Endrin 1.6 concentrate using 1 1/2 quarts per 100 gallons of water. **MEADOW MOUSE** (*microtus sp.*) control. Saturate ground cover under trees. Dilute Endrin 1.6 concentrate using one quart per 100 gallons of water.

SA BRAND 50[®]

1.6 Endrin Emulsible Concentrate

(Contains 1.6 lbs. Endrin per Gallon)

For Pine & Meadow Mice Control in Apple Orchards

ACTIVE INGREDIENTS:

Endrin (hexachloroepoxyoctahydro-endo, endo-dimethanonaphthalene)

Xylene

INERT INGREDIENTS

19.0%

~~70.0%~~

11.0%

100.0%

BOOM APPLICATION. Apply 100 gallons of diluted spray in a band 11 feet wide and 670 feet long on each side of tree row (350-400 gallons per acre).

HAND GUN APPLICATION (power sprayer). Apply 11 gallons of spray under each mature tree (proportionately less for young trees). Treat from base of tree to one or two feet beyond the drip line.

Do not apply more than 1 1/2 gallons of 1.6 Endrin Concentrate or 400 gallons of diluted spray per acre by either hand gun or boom application.

Post and otherwise prevent entry to treated areas for 30 days following treatment. Do not graze orchards, cut forage for hay, or allow drop fruit to be used for any purpose.

Workers should not re enter treated area without protective equipment.

DO NOT USE POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

Decontaminate used containers by washing thoroughly with lye solution. Destroy and bury used containers. Do not reuse. Store in a safe place out of reach of children. Protect from temperatures below freezing.

This material may not give satisfactory control under all conditions. Severity of infestation, ground cover, weather conditions and method of application are beyond the control of the seller.


Buyer assumes all risks of use, storage and handling of this material not in strict accordance with the directions and instructions given herewith.

(See Side Panels for Instructions and Precautions in Using)

Southern Agricultural Insecticides, Inc.

Palmeto, Florida Boone, N. C. Hendersonville, N. C.

NET LIQUID CONTENT



DANGER! POISONOUS IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN. RAPIDLY ABSORBED THROUGH SKIN. Do not get in eyes, on skin, or on clothing. Do not breathe vapor or spray mist. Wash thoroughly with soap and water after handling and before eating or smoking. Wear clean clothing daily, clean synthetic rubber gloves and a mask or respirator of a type passed by the United States Bureau of Mines for Endrin protection. Shower and wash thoroughly and change to clean clothing after each day's spraying. Do not apply or allow drift to areas occupied by unprotected humans or beneficial animals. Do not contaminate feed or foodstuffs.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Not for use or storage in or around the home.

MAKE POSTHARVEST APPLICATION ONLY. Make single application to the orchard floor in October or November. Specific directions concerning nozzles should be obtained from State Agricultural Extension Specialist. Do not disturb trail system. Allow at least 2 months before and after treatment before cultivating or mowing.

PINE MOUSE (*pitymys pinetorum*) control. Spray must be applied at 500 lbs. per square inch pressure in order to penetrate ground cover to reach trail systems. Dilute Endrin 1.6 concentrate using 1½ quarts per 100 gallons of water. **MEADOW MOUSE** (*microtus sp.*) control. Saturate ground cover under trees. Dilute Endrin 1.6 concentrate using one quart per 100 gallons of water.

SA BRAND 50[®]

1.6 Endrin Emulsible Concentrate

(Contains 1.6 lbs. Endrin per Gallon)
For Pine & Meadow Mice Control in Apple Orchards

ACTIVE INGREDIENTS:
 Endrin (hexachloroepoxyoctahydro-endo, endo-dimethanonaphthalene)

Xylene
INERT INGREDIENTS

19.0%
~~70.0%~~
~~11.0%~~ 31%
 100.0%

 KEEP OUT OF REACH OF CHILDREN 
DANGER! POISON!

ANTIDOTE AND FIRST AID TREATMENT
 GASTROINTESTINAL: In case of suspected or confirmed poisoning, induce vomiting if the victim is conscious. If the victim has been swallowed and is unconscious, immediately call a physician and give warm milk or water. If the victim is unconscious, do not wait for a physician. Turn the victim onto his side. Remove all vomit. If the victim is unconscious, do not give anything by mouth. Do not induce vomiting. Keep patient quiet and quiet.
 EXTERNAL: In case of contact, immediately remove contaminated clothing and wash the skin with soap and running water. If not, get the victim to a hospital immediately. Wash the skin with soap and water for 15 minutes and then treat as usual.
 (See Side Panels for Instructions and Precautions in Using)

Southern Agricultural Insecticides, Inc.
 Palmeto, Florida Boone, N. C. Hendersonville, N. C.

BOOM APPLICATION. Apply 100 gallons of diluted spray in a band 11 feet wide and 670 feet long on each side of tree row (350-400 gallons per acre).

HAND GUN APPLICATION (power sprayer). Apply 11 gallons of spray under each mature tree (proportionately less for young trees). Treat from base of tree to one or two feet beyond the drip line.

Do not apply more than 1½ gallons of 1.6 Endrin Concentrate or 400 gallons of diluted spray per acre by either hand gun or boom application.

Post and otherwise prevent entry to treated areas for 30 days following treatment. Do not graze orchards, cut forage for hay, or allow drop fruit to be used for any purpose. Workers should not re-enter treated area without protective equipment.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

Decontaminate used containers by washing thoroughly with lye solution. Destroy and bury used containers. Do not re-use. Store in a safe place out of reach of children. Protect from temperatures below freezing.

NEED TO PHYSICIAN in case of poisoning. For information on the use of this product, contact the manufacturer, Southern Agricultural Insecticides, Inc., Palmeto, Florida. For information on the use of this product, contact the manufacturer, Southern Agricultural Insecticides, Inc., Palmeto, Florida.

This material may not give satisfactory control under all conditions. Severity of infestation, ground cover, weather conditions and method of application are beyond the control of the seller.

Buyer assumes all risks of use, storage and handling of this material not in strict accordance with the directions and precautions given herewith.

EPA Reg. No. 829-98
NET LIQUID CONTENT