



Use as is, to control Mealybugs, Red Spider Mites, Aphids, Thrip (exposed), Whiteflies and Spittlebugs. Apply in a manner to provide complete and uniform coverage of infested plants. Be sure to get good coverage of underside of leaves and trunk or main stem. Avoid overspraying to the point of excessive runoff. Treat when pests appear and repeat at 7 to 10 day intervals, if needed.

NOTE: Do not use on Azaleas, Camellias, Poinsettias, Rose Bushes, or variegated Ivy because of possible injury to these plants.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.





53% EMULSIBLE MALATHION

ACTIVE INGREDIENTS:	
Malathion (0,0-Dimethyl phosphorodithioate of diethyl mercaptosuccinate).....	53.0%
INERT INGREDIENTS*.....	47.0%
TOTAL.....	100.0%

*Contains xylene range aromatic solvent.
Contains 5.0 lbs. Malathion per gallon.

Keep Out of Reach of Children CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Get medical attention immediately. Do not induce vomiting without advice of physician. Contains aromatic petroleum solvent. Cautious gastric lavage may be indicated.

IF IN EYES: Flush eyes with water for 15 minutes. Get medical attention.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration. Get medical attention.

IF ON SKIN: Immediately flush skin with soap and water for at least 15 minutes while removing contaminated clothing and shoes. Get medical attention.

NOTE TO PHYSICIAN: This product may cause cholinesterase inhibition. Atropine is antidotal. 2-PAM/Proto PAM may be effective, but only as an adjunct to atropine.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA EST NOP 829-FL-1 (3), 829-NC-1 (4), 829-NC-2 (5) Number following Est. No. corresponds to first digit of Lot No. on side of label.

Southern Agricultural Insecticides, Inc.
Palmetto, FL 34220 Hendersonville, NC 28793 Boone, NC 28607
EPA Reg. No. 829-101 J2 Net Contents Liquid: 1 Gallon

ANIMAL QUARTERS: For control of fleas under houses, apply a spray containing 10% of water at rate of 1 gallon per 1000 sq. ft. of manure and debris before spraying.

BEANS: For control of Mexican Bean Beetle when insects first appear. Use 2 teaspoonfuls of water per acre). Do not use within 1 day of harvest.

CITRUS: (Florida) for the control of purple scale use 2 teaspoonfuls per gallon of water (2 pints per gallon of water) spray post bloom and second during June. Use 5 gals. of 53% Malathion per acre in any spray.

CUCUMBERS and SQUASH: (In home gardens only), for the control of aphids and squash bugs when aphids first appear and repeat application when leaves appear or when plants are set in the field. Apply to obtain thorough coverage. Use 3 pints per 100 gals. of water per acre). Do not apply within 10 days of harvest. Do not apply to cabbage, collards, etc.

BROCCOLI, CABBAGE, COLLARDS, KALE: (In home gardens only), for the control of aphids and squash bugs when aphids first appear and repeat application when leaves appear or when plants are set in the field. Apply to obtain thorough coverage. Use 3 pints per 100 gals. of water per acre). Do not apply within 10 days of harvest. Do not apply to cabbage, collards, etc.

HOUSEHOLD INSECTS: For the control of cockroaches, clovermites, and earwigs in and about the house. Apply by means of a coarse spray nozzle. Apply under drain boards, under sinks, under stoves, and in other places where insects are likely to hide. A spray can cause staining to some finished surfaces.

Use only with adequate ventilation. After spraying, ventilate thoroughly before occupying enclosed spaces until sprays have dried.

STORAGE AND DISPOSAL:
Do not contaminate food or feed by storage or handling.
PESTICIDE STORAGE: Store pesticide in original container in a locked storage area.
PESTICIDE DISPOSAL: Securely wrap original container in paper and discard in trash.
CONTAINER DISPOSAL: Do not reuse or burn. Recycle if possible.
ing in trash.

Buyer assumes all risks of use, storage or handling in accordance with the directions given herewith.



in accordance with
Phase 2
Change in Status.