

LAWNS AND TURF AREAS NOT TO BE GRAZED

Do not exceed label directions. Shake well before using. Do not use on lawns or other turf areas which are to be grazed by animals. Do not apply to lawns which are to be grazed by animals. Do not apply to lawns which are to be grazed by animals.

This product is toxic to humans. Keep out of any body of water. Do not use on treated areas until dry.

ANTS: Apply 1/2 ounce per 100 sq ft of water to cover 100 sq ft. Repeat application to ant hills. Sprinkle over application. Repeat if necessary.

SOD WEBWORMS, ARMYWORMS, FALL ARMYWORMS, and MOLE CRICKETS: Spray with 2 tablespoons in 1/2 gallon of water to cover 100 sq ft. Apply as soon as infestation is noted. Worm control is more effective when worms are small. Repeat applications when necessary. For sod webworms, apply in late spring and early fall.

JAPANESE BEETLE (LARVAE): Spray with 1/2 tablespoonful in 1/2 gallon of water to cover 100 sq ft. Apply preferably in cool of early evening when exhumation of pupae shows. Do not apply when pupae are in soil. Water well into soil. One application should last several years.

CHIGGERS: Apply 1/2 ounce per 100 sq ft of lawn. Repeat application when necessary.

S A BRAND 5 0 0 50% CHLORDANE EMULSION CONCENTRATE

Contains 4 lbs. Technical Chlordane per gallon

Table with 2 columns: Ingredient, Amount. ACTIVE INGREDIENTS: Technical Chlordane 50.0, Petroleum Distillate 40.0. INERT INGREDIENTS: 100.0

*Equivalent to 50% DDT, 4, 7 - Methanorahydroindane and 70% related compounds.

KEEP OUT OF REACH OF CHILDREN

CAUTION: Harmful if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing spray mist. Avoid contact with skin, eyes and clothing. In case of contact with skin wash with soap and water. For eyes flush with water and get medical attention. Avoid contamination of food and foodstuffs. Do not use or store near heat or open flame.

ANTIDOTE:—Internal: Emetic of 1 tablespoonful of mustard in tumbler of water. Call a physician.

Southern Agricultural Insecticides, Inc. P.O. Box 111, Kinston, N.C. Hendersonville, N.C.

U.S. PAT. 2,811,111. N. C. PAT. 1,151,111.

HOUSEHOLD INSECTS

Do not allow children or animals to come in contact with treated surfaces until treated areas have completely dried. Do not treat without good ventilation. Spraying should be made so as to avoid fabrics. Chlordane emulsions can stain some surfaces. It is advisable to make a spot test spraying to determine stain resistance of surfaces to be treated.

ANTS, ROACHES, SILVERFISH: Dip a 1/2 cup-spoonful in a gallon of water. Apply with a paint brush or coarse wet spray. Thoroughly treat all cracks, baseboards, around plumbing, areas behind shelving, storage areas, around door sills, windows, screens and any other places where insects might hide or have entry. Repeat applications when necessary. Some roaches are resistant to chlordane treatments.

WASPS: Spray nest after dark when insects have returned to nest, paying particular attention to nest opening at rate of 1/2 tablespoonful per gallon of water.

SPIDERS AND CENTIPEDES: In basements and garages. Use 1/2 table-spoonful per gallon of water. Thoroughly spray or brush for cracks, pipes, pipes, storage areas, and other infested areas. Repeat applications when necessary.

Apply to any surface which is to be treated. Do not use this material on any treated surface with which it comes in contact.

LAWNS AND TURF AREAS NOT TO BE GRAZED

Do not exceed label directions. Shake well before using. Do not use on lawns or other turf areas which are to be grazed by animals. Do not apply to lawns which are to be grazed by animals. Do not apply to lawns which are to be grazed by animals.

ANTS: Apply 1/2 ounce per 100 sq ft of water to cover 100 sq ft. Repeat application to ant hills. Sprinkle over application. Repeat if necessary.

SOD WEBWORMS, ARMYWORMS, FALL ARMYWORMS, and MOLE CRICKETS: Spray with 2 tablespoons in 1/2 gallon of water to cover 100 sq ft. Apply as soon as infestation is noted. Worm control is more effective when worms are small. Repeat applications when necessary. For sod webworms, apply in late spring and early fall.

JAPANESE BEETLE (LARVAE): Spray with 1/2 tablespoonful in 1/2 gallon of water to cover 100 sq ft. Apply preferably in cool of early evening when exhumation of pupae shows. Do not apply when pupae are in soil. Water well into soil. One application should last several years.

CHIGGERS: Apply 1/2 ounce per 100 sq ft of lawn. Repeat application when necessary.

S A BRAND 5 0 0 50% CHLORDANE EMULSION CONCENTRATE

Contains 4 lbs. Technical Chlordane per gallon

Table with 2 columns: Ingredient, Amount. ACTIVE INGREDIENTS: Technical Chlordane 50.0, Petroleum Distillate 40.0. INERT INGREDIENTS: 100.0

*Equivalent to 50% DDT, 4, 7 - Methanorahydroindane and 70% related compounds.

KEEP OUT OF REACH OF CHILDREN

CAUTION: Harmful if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing spray mist. Avoid contact with skin, eyes and clothing. In case of contact with skin wash with soap and water. For eyes flush with water and get medical attention. Avoid contamination of food and foodstuffs. Do not use or store near heat or open flame.

ANTIDOTE:—Internal: Emetic of 1 tablespoonful of mustard in tumbler of water. Call a physician.

Southern Agricultural Insecticides, Inc. P.O. Box 111, Kinston, N.C. Hendersonville, N.C. U.S. PAT. 2,811,111. N. C. PAT. 1,151,111.

HOUSEHOLD INSECTS

Do not allow children or animals to come in contact with treated surfaces until treated areas have completely dried. Do not treat without good ventilation. Spraying should be made so as to avoid fabrics. Chlordane emulsions can stain some surfaces. It is advisable to make a spot test spraying to determine stain resistance of surfaces to be treated.

ANTS, ROACHES, SILVERFISH: Dip a 1/2 cup-spoonful in a gallon of water. Apply with a paint brush or coarse wet spray. Thoroughly treat all cracks, baseboards, around plumbing, areas behind shelving, storage areas, around door sills, windows, screens and any other places where insects might hide or have entry. Repeat applications when necessary. Some roaches are resistant to chlordane treatments.

WASPS: Spray nest after dark when insects have returned to nest, paying particular attention to nest opening at rate of 1/2 tablespoonful per gallon of water.

SPIDERS AND CENTIPEDES: In basements and garages. Use 1/2 table-spoonful per gallon of water. Thoroughly spray or brush for cracks, pipes, pipes, storage areas, and other infested areas. Repeat applications when necessary.