

BRAND


**ACTIVE INGREDIENTS:** Carbaryl (1-Naphthyl N-methylcarbamate\*) INERT INGREDIENTS .....

\*Trademark of Union Carbide Corporation for: 1-Naphthyl N-methylcarbamate

100.00%

\*U. S. Pat. No. 2,903,478

## Keep Out Of Reach Of Children



May be used to day of harvest on listed crops except as otherwise indicated.

Where dosage ranges are indicated, use higher dosage on maturing plants and proportionatly less when applied by ground equipment to smaller plants. Do not exceed maximum indicated dosages. Before applying consult proper state extension authority to determine if local recommendations are consistant with this label. This product may injure tender plant parts particularly when applications are uneven, or when plants are wet, or when applications are made during periods or when plants are wet, or when applications are made curing periods of sustained high humidity. This product may not give satisfactory control under all conditions. Timing, method of application, weather conditions, severity of insect infestation and other factors influencing the results obtained from the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with the directions and precautions given herewith.

## Southern Agricultural Insecticides, Inc.

PALMETTO, FLA. BOONE, N. C.

HENDERSONVILLE, N. C.

U. S. D. A. REG. NO. 829-128

NET WEIGHT 4 LBS.

Beans and cowpeas for the control of Mexican bean beetle, Japanese beetle adults, bean leaf beetle, cucumber beetle, corn ear worm, and fall army worm. Make 20 to 40 lbs. per acre applications on beans and 20 to 30 lbs. per acre applications on cowpeas. Apply to foliage as needed to maintain control.

Alfalfa, Clover, Grass and Hay, and Soybeans to control Mexican bean beetle, Japanese beetle adults, army worms, leafhoppers, grasshoppers, corn ear worm, velvet bean caterpillars, make 30 lbs, per acre applica-

Broccoli, Cabbage, and Cauliflower for the control of cabbage caterpillars and harelquin bug, apply at weekly intervals using 20 to 40 lbs. per acre applications depending on the size of the plants, beginning when plants are set in the field. Control of cabbage loopers with this dust may not be satisfactory. Do not apply to cabbage, broccoli, or

caulillower within 3 days of harvest. Corn for the control of corn ear worms, fall army worms, sap beetles, Japanese beetle adults. Corn ear worms, tan army worms, sap peetles, Japanese beetle adults. Corn ear worm and fall army worm larvae and sap bug in silks: Make four 40 lbs. per acre applications, directed to ear and silks at 2 day intervals, beginning when silks first appear. Continue applications at 3 day intervals until silks have completely wilted. Corn ear worm and fall army worm larvae in the whorl (bud worm damage). When insects appear direct dust into the whorl using 30 to 40 lbs. per acre applications. Repeat if necessary, Japanese beetle adults: Make 40 lbs. per acre applications, as necessary to maintain control. Satisfactory worm control in corp is diffigult to obtain, consult the proper State Extension: Authority for recommended frequency and timing of applications. Application of this product to the tassel region of corn during the pollen shed period will seriously

reduce bee populations.

Tomatoes, Potatoes, Pepper, Okra and Eggplant for the control of Colorado potato beetleflea bettles, leafhoppers, tomato fruit worm, fall army worm, tomato horn worm, and stink bugs. Make 25 to 40 lbs. per acre applications to foilage as necessary. For tomato fruit worm, on tomatoes, peppers, and okra apply at weekly intervals after fruit has begun to set.

Grapes for the control of Japanese beetle adults, apply 40 lbs. per acre when insects first appear, repeat as necessary. For control of grape berry moth, flea beetle, leafhoppers, and leaf skeletonizer con-

sult proper State Extension authority for timing and frequency of

applications.

Chickens Northern fowl mite and lice and as a supplement to premise treatment for chicken mites. Dust birds thoroughly using 1 lb. per treatment for chicken mites. Dust birds thoroughly using 1 lb. per 100 birds. Repeat treatment in 4 weeks if needed, but not more often. Do not apply within 7 days of slaughter. Chicken houses for the control of chicken mites, apply 1 lb. per 40 sq. ft. of litter and roosts. Obtain thorough coverage. Force dust into all cracks and crevices. Do not apply directly to eggs or nest litter. Do not house birds in treated houses within 7 days of slaughter. Do not treat when birds are under influence of stress.

are under influence of stress.

Dogs and cats for the control of brown dog tick and fleas, apply thoroughly, rubbing well into skin. Also dust animal sleeping quarters. Repeat at weekly intervals as needed. Do not treat nursing pupples, kittens under 4 weeks of age, or animals in a weakened condition.

NOTE: When treating animal premises do not contaminate feed, water or utensile.

water or utensils.

Lawns and Turi for the control of sod webworms and fall armyworms when insects appear, apply evenly, using 5 lbs. per 1000 sq. ft. in late afternoon. Repeat if necessary.

Ornamentals in home garden: Arbor vitae, azalea, birch, carnation, chrysanthemum, dogwood elm, gladoli, hydrangea, juniper, lilac, maple, oak, rose, zinnia, to control bag worm, birch leaf miner, blister beetle, Japanese beetle and June beetle adults, lace bug, leaf rollers, mimosa webworm, orange tortrix, periodical cicada, tent caterpillar, thorn bug, and exposed thrips. Apply to plants as needed to maintain control. Consult state extension specialist for timing of applications. Do not use on Boston Ivy.

U. S. D. A. REG. NO. 829-128



## ACCEPTED

MAY 6 1971 FUNGICIDE AND ROO NTICIDE ACT 

829-128

## ACCEPTED

829- 128