


(A) 	United States Environmental Protection Agency Office of Pesticide Programs (H7505C) Washington, DC 20460	<input type="checkbox"/> Registration <input checked="" type="checkbox"/> Amendment <input type="checkbox"/> Other	OPP Identifier Number 189415
	Application for Pesticide:		

Section I

1. Company/Product Number 829-128	2. EPA Product Manager Dennis Edwards	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) SA-50 Sevin Brand Carbaryl Insecticide 5% Dust	PM# 19	
5. Name and Address of Applicant (Include ZIP Code) Southern Agricultural Insecticides, Inc. P.O. Box 218 Palmetto, FL 34420 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section II

<input type="checkbox"/> Amendment - Explain below	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - explain below.


Explanation: Use additional page(s) if necessary. (For section I and Section II.)
Label Change in accordance with PR notice 93-3

NOTIFICATION LABEL NOT REVIEWED PER PR NOTICE 93-3
DATE 7/1/93

Section III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
				<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
* Certification must be submitted.	If "Yes," Unit Package wgt.	No. per container	If "Yes," Package wgt.	No. per container	<input type="checkbox"/> Other (Specify) _____
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container	4. Size(s) of Retail Container	5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product			
6. Manner in Which Label Is Affixed To Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other (_____)		

Section IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Ronald L. Costa Rica	Title Registration Specialist	Telephone No. (Include Area Code) 813-722-3285
2. Signature 		6. Date Application Received (Stamped)
3. Title Registration Specialist		
4. Typed Name Ronald L. Costa Rica		
5. Date		

BEST AVAILABLE COPY



SEVIN®

BRAND CARBARYL INSECTICIDE

5% DUST

ACTIVE INGREDIENTS:

Carbaryl (1-Naphthyl N-methylcarbamate) 5.00%

INERT INGREDIENTS 95.00%

TOTAL 100.00%

SEVIN® is a registered trademark of Rhone-Poulenc for Carbaryl Insecticide

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician immediately. Induce vomiting immediately. To induce vomiting give one or two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.
IF INHALED: Remove victim to fresh air. Apply respiration if indicated.
IF ON SKIN: Wash promptly with soap and water. Rinse thoroughly.
IF IN EYES: Flush eyes thoroughly with water for at least 15 minutes. Call a physician immediately.
NOTE TO PHYSICIAN: Atropine sulfate is antidotal.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic and estuarine invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not use this product in commercial food areas or food handling establishments, restaurants, or other places where food is prepared or processed.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

EPA EST. NO. 829-FL-1 (3); 829-NC-1 (4); 829-NC-2 (5) Number following Est. No. corresponds to first digit in Lot No. on bottom of bag

Southern Agricultural Insecticides, Inc.

Palmetto, FL 34220

Hendersonville, NC 28703

Bohler, NC 28607

EPA REG. NO. 829-128

J2

Net Weight: 25 lbs.

BEST AVAILABLE COPY

NOTIFICATION
LABEL NOT REVIEWED
PER PR NOTICE 9-3
DATE 7/1/93

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

NOTE: May be used to day of harvest on listed crops except as otherwise indicated. Where dosage ranges are indicated, use higher dosages on maturing plants and proportionately less when applied by ground equipment to smaller plants. Do not exceed maximum indicated dosages. Before applying consult proper State Extension Authority to determine if local recommendations are consistent with this label. This product may injure tender plant parts particularly when applications are uneven, or when plants are wet, or when applications are made during periods of sustained high humidity. This product may not give satisfactory control under all conditions. Timing, method of application, weather conditions, severity of insect infestation and other factors influencing the results obtained from the use of this product are beyond the control of the seller.

BEANS and COWPEAS for the control of Mexican bean beetle, Japanese beetle adults, bean leaf beetle, cucumber beetle, corn earworm, and fall armyworm. Make 20 to 40 lbs. per acre applications on beans and 20 to 30 lbs. per acre applications on cowpeas. Apply to foliage as needed to maintain control. Do not apply to cowpeas within 3 days of harvest.

ALFALFA, CLOVER, GRASS and HAY and SOYBEANS to control Mexican bean beetle, Japanese beetle adults, armyworms, leafhoppers, grasshoppers, corn earworm, velvet bean caterpillars. Make 30 lbs. per acre applications as needed. Do not apply to alfalfa within 3 days of harvest.

BROCCOLI, CABBAGE and CAULIFLOWER for the control of cabbage caterpillars or harlequin bug. Apply at weekly intervals using 20 to 40 lbs. per acre applications depending on the size of the plants, beginning when plants are set in the field. Control of cabbage loopers, with this dust may not be satisfactory. Do not apply to cabbage, broccoli, or cauliflower within 3 days of harvest.

CORN for the control of corn earworms, fall armyworms, sap beetles, Japanese beetle adults, corn earworm and fall armyworm larvae in silks: Make four 40 lbs. per acre applications, directed to ear and silks at 2-day intervals beginning when silks first appear. Continue applications at 3-day intervals until silks have completely withed. Corn earworm and fall armyworm larvae in the whorl (bud worm damage): When insects appear direct dust into the whorl using 30 to 40 lbs. per acre application. Repeat if necessary. Japanese beetle adults: Make 40 lbs. per acre applications as necessary to maintain control. Satisfactory worm control in corn is difficult to obtain. Consult the proper State Extension Authority for recommended timing of applications. Application of this product to the tassel region of corn during pollen shed period will seriously reduce bee population.

TOMATOES, POTATOES, PEPPER, OKRA and EGGPLANT for the control of Colorado potato beetle, flea beetle, leafhoppers, tomato fruitworm, fall armyworm, tomato hornworm and stink bugs. Make 25 to 40 lbs. per acre applications to foliage as necessary. For tomato fruitworm on tomatoes, peppers and okra apply at weekly intervals after fruit has begun to set.

GRAPES for the control of Japanese beetle adults, apply 40 lbs. per acre when insects first appear. Repeat as necessary. For control of grape berry moth, flea beetle, leafhoppers, and leaf skeletonizer consult proper State Extension authority for timing of applications.

CHICKENS: Northern lowl mites and lice and as a supplement to premise treatment for chicken mites. Dust birds thoroughly using 1 lb. per 100 birds. Repeat treatment in 4 weeks if needed, but not more often. Do not apply within 7 days of slaughter. Chicken houses for the control of chicken mites, apply 1 lb. per 40 sq. ft. of litter and roosts. Obtain thorough coverage. Force dust into all cracks and crevices. Do not apply directly to eggs or nest litter. Do not house birds in treated houses within 7 days of slaughter. Do not treat when birds are under influence of stress.

DOGS and CATS for the control of brown dog ticks and fleas. Dust liberally over animal rubbing well into skin. Do not apply to areas around eyes, nose and mouth. Prevent dust from entering animals nose and mouth. Also dust animal sleeping quarters. Repeat at weekly intervals as needed. Do not use on pregnant dogs, nursing puppies, kittens under 4 weeks of age, or animals in a weakened condition.

NOTE: This product may stain carpets or other fabrics. Do not apply directly to carpets or allow the treated animals on carpeted areas.

LAWN and TURF for the control of sod webworms and fall armyworms. When insects appear, apply evenly, using 5 lbs. per 1000 sq. ft. in late afternoon. Repeat if necessary.

ORNAMENTALS in home gardens. Use on arborvitae, azalea, birch, carnation, chrysanthemum, dogwood, elm, gladiol, hydrangea, juniper, lilac, maple, oak, rose, zinnia to control bagworm, birch leafminer, blister beetle, Japanese beetle and June beetle adults, lace bug, leaf rollers, mimosa webworm, orange tortrix, periodical cicada, tent caterpillar, thorn bug, and exposed thrips. Apply to plants as needed to maintain control. Consult State Extension specialist for timing of applications. Do not use on Boston ivy.


NOTE: Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s).

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal. **STORAGE:** Store pesticide in the original container in a locked storage area. **PESTICIDE DISPOSAL:** To dispose of excess pesticide, securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with the directions and precautions given herewith

EPA REG. NO. 829-128



0 51538 10647 4

BEST AVAILABLE COPY