

AL...  
 JUN 3, 71  
 024-124

Finished label out size: 6 1/2" x 10"

**CAUTION**

Do not apply SA-50 2,4-D AMINE WEED KILLER directly to, or otherwise permit it to come into contact with vegetables, flowers, grapes, fruit trees, ornamentals, cotton, or other desirable plants which are sensitive to 2,4-D and similar materials, and do not permit spray mist containing 2,4-D to drift onto them, since even minute quantities of the spray may cause severe injury during both growing and dormant periods. Coarse sprays are less likely to drift. Do not contaminate irrigation ditches or water used for domestic purposes. Do not store near fertilizers, seeds, insecticides, or fungicides. Excessive amounts of 2,4-D in the soil may temporarily inhibit seed germination or plant growth. To avoid injury to desirable plants do not store, handle or apply other agricultural chemicals with the same containers or equipment used for 2,4-D.

Local conditions may affect the use of herbicides. State agricultural authorities in many states issue recommendations to fit local conditions.

**NOTE**

Mix SA-50 2,4-D AMINE WEED KILLER only with water. Use only as suggested. In mixing sprays, add half the water to the tank, then add the SA-50 2,4-D AMINE WEED KILLER with agitation, then add the balance of water with continued agitation. Apply at low pressure during warm weather when weeds are young and growing actively.

**Lawn Grasses**

For use on Bermuda, Marion Blue, Kentucky Blue, Red Top Fescue, and Zoysia. Do not use on Dichondra, St. Augustine, Carpet Grass, Centipede, Bent, or other creeping grasses, or on putting greens. Most legumes are easily damaged. Do not apply to turf that is under stress from nematode infestation, drought, or other cause.

Use 3 pints per acre or 6 teaspoonsful per 1000 square feet. Do not use on newly planted or seeded turf until grass has become well established.

**SA-50® BRAND**

**2,4-D AMINE WEED KILLER**

Contains 4 Lbs. per Gallon Acid Equivalent

**ACTIVE INGREDIENTS:**

Alkanolamine Salts (of the Ethanol and Isopropanol series) of 2,4-Dichlorophenoxyacetic Acid 59.0%

2,4-Dichlorophenoxyacetic Acid Equivalent 38.0  
 INERT INGREDIENTS 41.0%

100.0%

**Keep Out of Reach of Children**

**CAUTION!**

Harmful if swallowed.

Causes irritation of skin and eyes. Do not get in eyes. Avoid contact with skin and clothing.

In case of contact with eyes, flush with plenty of water for at least 15 minutes and get medical attention; wash skin with soap and plenty of water. Remove and wash contaminated clothing before reuse. Do not wear contaminated shoes.

SEE SIDE PANELS FOR ADDITIONAL CAUTIONS AND DIRECTIONS. READ ENTIRE LABEL BEFORE USING.

Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions and precautions given hereon.

NET CONTENTS: ONE QUART LIQUID

**SOUTHERN AGRICULTURAL INSECTICIDES, INC.**

GREENSBORO, N. C. BOONE, N. C. PALMETTO, FLA.

**Corn—Pre-Emergency Treatment**

Apply evenly to the soil surface 3 to 5 days after planting corn but before corn emerges. Use lower rate on light soils; higher rate on heavy soils. Do not use SA-50 2,4-D AMINE WEED KILLER on sandy soils. Mix 1/2 cup to 1 cup of SA-50 2,4-D AMINE WEED KILLER in 3 gallons of water to cover 1/10 acre, or the contents of this container in 7 1/2 to 10 gallons of water to cover 1/2 to 1 acre.

**Corn—Post Emergence**

Apply evenly to soil surface when corn is 10 to 12 inches tall. Direct spray so as to avoid getting spray into the whorl of corn or on corn leaves. Mix 3 to 4 tablespoonsful of SA-50 2,4-D AMINE WEED KILLER in 3 gallons of water to cover 1/10 acre, or mix the contents of this container in 60 gallons of water to cover 2 acres. Do not cultivate for at least 10 days after spraying. Application of SA-50 2,4-D AMINE WEED KILLER to wet soils during hot weather may cause injury to corn.

**General Weed Control**

In vacant lots, on roadsides, and along fence rows, use 3 tablespoonsful of SA-50 2,4-D AMINE WEED KILLER per gallon of water or 3 quarts per 100 gallons of water, applying enough to wet all foliage. Deep-rooted perennial weeds may require repeated applications as new growth appears.

SA-50 2,4-D AMINE WEED KILLER is recommended for control of the following, in addition to many other broad-leaved weeds, without injury to most grasses:

- |                 |                    |
|-----------------|--------------------|
| alligator weed  | pigweed            |
| dandelion       | plantain           |
| docks           | ragweed            |
| dog fennel      | shepherd's-purse   |
| ranweed         | tarweed            |
| goldenrod       | thistles           |
| krackia         | vetch              |
| lamb's quarters | wild carrot        |
| mustards        | wild morning glory |
| poppygrass      | wild radish        |

Always use a spray empty container with care.

**ACCEPTED**  
**JUN 3, 71**  
 UNDER THE FEDERAL INSECTICIDE  
 FUNGICIDE AND ROENTGENIC ACT  
 FOR ECONOMIC PEST CONTROL  
 REGISTERED UNDER NO. 829-129 CT

**Finished label out size: 6 1/2" x 10"**

**CAUTION**

Do not apply SA-50 2,4-D AMINE WEED KILLER directly to, or otherwise permit it to come into contact with vegetables, flowers, grapes, fruit trees, ornamentals, cotton, or other desirable plants which are sensitive to 2,4-D and similar materials, and do not permit spray mist containing 2,4-D to drift onto them, since even minute quantities of the spray may cause severe injury during both growing and dormant periods. Coarse sprays are less likely to drift. Do not contaminate irrigation ditches or water used for domestic purposes. Do not store near fertilizers, seeds, insecticides, or fungicides. Excessive amounts of 2,4-D in the soil may temporarily inhibit seed germination or plant growth. To avoid injury to desirable plants do not store, handle or apply other agricultural chemicals with the same containers or equipment used for 2,4-D.

Local conditions may affect the use of herbicides. State agricultural authorities in many states issue recommendations to fit local conditions.

**NOTE**

Mix SA-50 2,4-D AMINE WEED KILLER only with water. Use only as suggested. In mixing sprays, add half the water to the tank, then add the SA-50 2,4-D AMINE WEED KILLER with agitation, then add the balance of water with continued agitation. Apply at low pressure during warm weather when weeds are young and growing actively.

**Lawn Grasses**

For use on Bermuda, Marion Blue, Kentucky Blue, Red Top Fescue, and Zoysia. Do not use on Dichondra, St. Augustine, Carpet Grass, Centipede, Bent, or other creeping grasses, or on putting greens. Most legumes are easily damaged. Do not apply to turf that is under stress from nematode infestation, drought, or other causes.

Use 3 pints per acre or 6 teaspoonsful per gallon per 1000 square feet. Do not use on newly planted or seeded turf until grass has become well established.

U. S. Patent Numbers:  
 2,800,941; 2,896,518; 2,458,983; 2,515,198;  
 Re 28; 116

**SA-50® BRAND**

**2,4-D AMINE  
 WEED KILLER**

**Contains 4 Lbs. per Gallon Acid Equivalent**

**ACTIVE INGREDIENTS:**

Alkanolamine Salts (of the Ethanol and Isopropanol series) of 2,4-Dichlorophenoxyacetic Acid

2,4-Dichlorophenoxyacetic Acid Equivalent 38.0

**INERT INGREDIENTS**

**Keep Out of Reach of Children  
 CAUTION!**

**Harmful if swallowed.**

**Causes irritation of skin and eyes. Do not get in eyes. Avoid contact with skin and clothing.**

In case of contact with eyes, flush with water for at least 15 minutes and get medical attention; wash skin with soap and plenty of water. Do not move and wash contaminated clothing before reuse. Do not wear contaminated shoes.

SEE SIDE PANELS FOR ADDITIONAL CAUTIONS AND DIRECTIONS. READ ENTIRE LABEL BEFORE USE.

Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions and cautions given herewith.

NET CONTENTS: ONE QUART LIQUID

**SOUTHERN AGRICULTURAL  
 INSECTICIDES, INC.**

HENDERSONVILLE, N. C. BOONE, N. C. PALM BEACH, FLA.  
 U. S. D. A. Reg. No. 829-129 B