

TRADE MARK AND REGISTERED SERVICE MARK
FOR SEVIN DUST, TRADE MARK
OF SYNGENTHA S.A.

SA BRAND 50[®] SEVIN^{*} 1.75% DUST

ACTIVE INGREDIENTS:

Carbaryl (1-Naphthyl N-methylcarbamate) 1.75%

INERT INGREDIENTS 98.25%

*Trade Mark of Froom Chemical Corporation, Fort Worth, Texas, U.S.A. Pat. Nos. 2,398,175
and 2,411,711

FOR USE ON

Beans to control Mexican bean beetle and bean leaf beetle. Apply 60 lbs. per acre to foliage as needed.

Cucumbers and Squash to control cucumber beetles. Apply 60 lbs. per acre to foliage as needed.

Potatoes and Tomatoes to control Colorado potato beetle. Apply 75 lbs. per acre to foliage as eggs begin to hatch. Repeat as needed.

Keep Out of Reach of Children

CAUTION! Harmful if swallowed or inhaled. May irritate eyes, nose, throat, and skin. Do not breathe dust. Avoid contact with skin, eyes, and clothing. Wear clean clothing daily. Wash thoroughly after using. Do not contaminate feed or foodstuffs.

Burn empty container. Stay out of smoke and fumes.

Note to Physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Do not contaminate any body of water nor allow drift from the crop area treated. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your cooperative Agricultural Extension Service.

May be used to day of harvest on listed crops except as otherwise indicated.

Dosages are indicated for mature plants. Use proportionately less when applied by ground equipment to smaller plants. Do not exceed maximum indicated dosages. Before applying consult proper technical or extension authority to determine if local recommendations are consistent with this label. This product may injure tender plant parts, particularly when applications are made on or when plants are wet, or when applications are made during periods of sustained high humidity. This product may not give satisfactory control under all conditions. Timing, method of application, weather conditions, severity of insect infestation and other factors influencing the result obtained from the use of this product are beyond the control of the manufacturer. Buyer assumes responsibility for proper handling of this material, not only in accordance with the directions, but in connection with other factors.

Southern Agricultural Insecticides, Inc.

REGISTERED TRADE MARK
© 1954 UNION CARBIDE CORPORATION
NEW YORK, N. Y.

SA BRAND 50[®] SEVIN* 1.75% DUST

ACTIVE INGREDIENTS:
Carbaryl (1-Naphthyl N-methylcarbamate) 1.75%
INERT INGREDIENTS 98.25%

*Trade Mark of Union Carbide Corporation for 1-Naphthyl N-methylcarbamate U. S. Pat. No. 2,903,478

FOR USE ON

Beans to control Mexican bean beetle and bean leaf beetle. Apply 60 lbs. per acre to foliage as needed.
Cucumbers and Squash to control cucumber beetles. Apply 60 lbs. per acre to foliage as needed.
Potatoes and Tomatoes to control Colorado potato beetle. Apply 75 lbs. per acre to foliage as eggs begin to hatch. Repeat as needed.

Keep Out of Reach of Children

CAUTION! Harmful if swallowed or inhaled. May irritate eyes, nose, throat, and skin. Do not breathe dust. Avoid contact with skin, eyes, and clothing. Wear clean clothing daily. Wash thoroughly after using. Do not contaminate feed or foodstuffs.

Burn empty container. Stay out of smoke and fumes.
Note to Physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Do not contaminate any body of water nor allow drift from the crop area treated. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your cooperative Agricultural Extension Service.

May be used to day of harvest on listed crops except as otherwise indicated.

Dosages are indicated for mature plants. Use proportionately less when applied by ground equipment to smaller plants. Do not exceed maximum indicated dosages. Before applying consult proper state extension authority to determine if local recommendations are consistent with this label. This product may injure tender plant parts, particularly when applications are made in the evening, or when plants are wet, or when applications are made during periods of sustained high humidity. This product may not give satisfactory control under all conditions. Timing, method of application, weather conditions, severity of insect infestation and other factors influencing the results obtained from the use of this product are beyond the control of the seller. Buyer assumes all risks of loss, damage, or handling of this material not in strict accordance with the directions and precautions given hereon.

Southern Agricultural Insecticides, Inc.

11000 W. 11th St., Tulsa, Okla. 74117
201-111