

DIRECTIONS

Mix SA-50 Maneb Spray Concentrate with water using ONE LEVEL TABLESPOONFUL PER GALLON OF SPRAY. Apply to obtain thorough coverage only. Make applications when plants are thoroughly dry and drying conditions are good. Apply at 5 to 7 day intervals and following rains. SA-50 Brand Maneb spray concentrate is compatible with most insecticides except spray oils and highly alkaline materials. Remove residue from sprayed vegetables by washing.

Do not reuse container. Destroy when empty.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes.

ROSES: For the control of black spot—begin applications when first leaf buds begin to unfold. Spray ground around plants generously to help control black spot infection from old leaves.

GLADIOLUS: To control Botrytis and Cercularia. Begin when first leaves appear. Continue sprays after cutting to maintain Botrytis control.

TOMATOES and POTATOES: For the control of early blight and late blight, gray leaf spot (Stemphylium) and anthracnose. For tomatoes begin when plants are set or first true leaves appear, for potatoes begin when plants are 2 to 6 inches tall. On tomatoes, do not apply within 5 days of harvest.

CUCUMBER and SQUASH: For the control of downy mildew and scab, begin applications when first true leaves appear. For scab control apply twice weekly after blooms begin to form. Do not apply within 5 days of harvest.

S A BRAND 5 0 0 Maneb Spray Concentrate

FOR THE CONTROL OF MANY PLANT DISEASES
On Roses—Tomatoes and Other Plants as indicated

ACTIVE INGREDIENTS:

Maneb (Manganese ethylenebisdithiocarbamate) 75.0%
(Manganese equivalent as Mn 15.4%)

INERT INGREDIENTS 25.0%

U.S. Pats. 2,504,404 - 2,710,882 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION: May be absorbed through skin and throat. Do not breathe dust. Keep away from fire and sparks. May cause irritation to eyes, nose, throat, and skin. Wash thoroughly after handling.

Southern Agricultural Insecticides, Inc.

Palmetto, Florida Boone, N.C. Hendersonville, N.C.
EPA Reg. No. 829-140 C Net Wt. 6 oz.

CABBAGE, CAULIFLOWER, and BROCCOLI: For the control of Alternaria Leaf Spot and Downy Mildew. Begin when plants are set in the field. Do not apply to cabbage, cauliflower, or broccoli within 7 days of harvest.

CANTALOUPE, HONEYDEW, MELONS AND WATERMELONS: For the control of Anthracnose, Downy Mildew, Leaf Spot, Downy Mildew and Downy Stem Blight. Begin applications at late bunching when vines begin to run or when disease first appears. Do not use within 5 days of harvest.

COLLARDS, KALE, MUSTARD GREENS AND TURNIPS: For control of Alternaria Leaf Spot and Downy Mildew. Begin applications when disease is first reported in the area. Do not apply within 10 days of harvest.

TOBACCO: For the control of Blue Mold in plant beds. Apply to obtain thorough coverage using 1 level teaspoonful per gallon of water. Spray only when plants are completely dry and drying conditions are good. Make first application when plants are about the size of a dime. Following applications twice per week until plants are free of Blue Mold and ready for transplanting. (Usually 8 to 12 applications.)

Follow directions carefully. Timing and method of applications, weather conditions, mixtures with other chemicals, and other influencing factors are beyond control of the seller. Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with the directions and precautions given herewith.

ACCEPTED

JAN 11 1973

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 829-140 SUBJECT
TO ATTACHED COMMENTS.