

# SA BRAND 50<sup>®</sup> Fruit Tree Spray

FOR THE CONTROL OF  
Many Fruit Insects and Diseases  
IN HOME PLANTINGS

**ACTIVE INGREDIENTS:**

Ferbam (Ferric dimethyl dithiocarbamate) 15.0%

Malathion (0 0-dimethyl dithiophosphate of diethyl mercaptosuccinate) 9.5%

Sulfur 23.6%

**INERT INGREDIENTS**

51.9%

100.0%

## Keep Out of Reach of Children

**CAUTION!** Harmful if inhaled or swallowed. Avoid breathing concentrate or spray mist. Avoid repeated or prolonged contact with skin, eyes, or clothing. Wash thoroughly after using. In case of contact with eyes, flush with plenty of water and get medical attention. Do not contaminate food and foodstuffs. Use empty containers away from smoke and fumes. Store in a safe, dry place.

Toxic to fish. Do not contaminate any part of water.

This product may not give satisfactory control under all conditions. Timing and method of applications, weather conditions, mixtures with other chemicals, and other influencing factors are beyond control of the seller. Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with the directions and precautions given hereon.

## Southern Agricultural Insecticides, Inc.

Palmetto, Florida Boone, N. C. Hendersonville, N. C.

U. S. E. A. REG. NO. 147-118-B

NET WEIGHT 3 LBS.

## FOR USE ON

**APPLES** for the control of scab, rust, powdery mildew, leaf spots, bitter rot, sooty blotch, aphids, leafroller, codling moth, Japanese beetle adults, oriental fruit moth, and spider mites. Apply first application when buds show 1/4 in. to 1/2 in. new growth; second application when center buds first show pink, until just before bloom; third application when most of the petals have dropped. Make four or five sprays at two-week intervals until one week before harvest. Do not apply to apples within seven days of harvest. Applications to McIntosh, Cortland, Golden, and related varieties may cause injury.

**GRAPES** to control black rot, leafhoppers, flea beetles, aphids, and Japanese beetle adults. Apply as follows: when new shoots are 1 to 2 inches long, when shoots are 7 to 8 inches long just before blossoms open, just after bloom, 10 days after bloom, and thereafter at two-week intervals. Do not apply closer than 7 days of harvest.

**RASPBERRIES AND DEWBERRIES** to control anthracnose, spur blight and rust. Also to control aphids and scale insects when these insects are present at the indicated times of application. Apply 1 1/2 lbs. in 10 gallons of water (9 tablespoonfuls per gallon) as a delayed dormant spray. Then apply 1 lb. in 10 gallons of water (6 tablespoonfuls per gallon) as follows: just before blossoms open, when new canes are 1 1/2 feet high, and just after harvest. Do not apply within 40 days of harvest.

For the above indicated sprays apply at the rate of 1 1/2 lbs. SA-50 Fruit Tree Spray to 10 gallons of water, or 6 lbs. or 12 tablespoonfuls per gallon of water, except as otherwise indicated. Apply to obtain thorough coverage. Applications should be made under good spraying conditions.

U. S. E. A. REG. NO. 147-118-B

**ACCEPTED**

FEB 6 1969

UNDER THE FEDERAL INSECTICIDE  
ACT AND RODENT CONTROL ACT  
AND UNDER THE FEDERAL POISON REGISTER  
ACT WITH NO RESERVATIONS SUBJECT  
TO REPEATED COMMENTS.