

SA BRAND 50®

FEB 7 1969

SEVIN* 3% DUST WITH 50% SULFUR

ACTIVE INGREDIENTS:	
Carbaryl (1-Naphthyl N-methylcarbamate)	3.00%
Sulfur	50.00%
INERT INGREDIENTS	47.00%

100.00%

*Trademark of Union Carbide Corporation for:
1-Naphthyl N-methylcarbamate U. S. Pat. No. 2,903,478

FOR USE ON beans and cowpeas for the control of bean leaf beetle, corn ear worm, cucumber beetle, Japanese beetle, Mexican bean beetle, potato leafhopper, and stink bugs. Apply to foliage as needed to maintain control using 65 lbs. per acre application on maturing plants. Use proportionately less depending on size of plants when applied by ground equipment to younger plants.

Eggplant for the control of Colorado potato beetle, flea beetles, and hornworms. Apply to foliage as needed to maintain control using 70 lbs. per acre application on maturing plants. Use proportionately less depending on the size of plants when applied by ground equipment to younger plants.

Do not exceed indicated dosages. There is no time limitation on the use of this dust for the indicated crops and dosages.

Keep Out of Reach of Children

CAUTION! Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust. Avoid skin contact. Wash thoroughly after using and before eating and smoking. Wear regular long sleeved work clothing. Change to clean clothing daily. Do not contaminate feed or foodstuffs.

NOTE TO PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal.

This product may cause plant injury particularly when applied when foliage is wet, humidity is high or when temperatures exceed 90° F. For protection of honeybees avoid use, if possible, when bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to move bees to a distance beyond the bee flight range until one week after application.

Do not contaminate any body of water. Avoid drift from the crop area treated.

This dust may not give satisfactory control under all conditions. Timing and method of application, weather and crop conditions, and other influencing factors are beyond the control of seller. Buyer assumes all risks of use, storage and handling of this product not in strict accordance with the directions and precautions given herewith.

Southern Agricultural Insecticides, Inc.

PALMETTO, FLA. HENDERSONVILLE, N. C. BOONE, N. C.

U. S. D. A. REG. NO. 829-145 B

NET WEIGHT _____

SA BRAND **50**[®]

FEB 7 1969

SEVIN* 3% DUST

WITH 50% SULFUR

ACTIVE INGREDIENTS:	
Carbaryl (1-Naphthyl N-methylcarbamate)	3.00%
Sulfur	50.00%
INERT INGREDIENTS	47.00%

*Trademark of Union Carbide Corporation for: **100.00%**
1-Naphthyl N-methylcarbamate U. S. Pat. No. 2,903,478

FOR USE ON beans and cowpeas for the control of bean leaf beetle, corn ear worm, cucumber beetle, Japanese beetle, Mexican bean beetle, potato leafhopper, and stink bugs. Apply to foliage as needed to maintain control using 65 lbs. per acre application on maturing plants. Use proportionately less depending on size of plants when applied by ground equipment to younger plants.

Eggplant for the control of Colorado potato beetle, flea beetles, and hornworms. Apply to foliage as needed to maintain control using 70 lbs. per acre application on maturing plants. Use proportionately less depending on the size of plants when applied by ground equipment to younger plants.

Do not exceed indicated dosages. There is no time limitation on the use of this dust for the indicated crops and dosages.

Keep Out of Reach of Children

CAUTION! Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust. Avoid skin contact. Wash thoroughly after using and before eating and smoking. Wear regular long sleeved work clothing. Change to clean clothing daily. Do not contaminate feed or foodstuffs.

NOTE TO PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal.

This product may cause plant injury particularly when applied when foliage is wet, humidity is high or when temperatures exceed 90° F. For protection of honeybees avoid use, if possible, when bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to move bees to a distance beyond the bee flight range until one week after application.

Do not contaminate any body of water. Avoid drift from the crop area treated.

This dust may not give satisfactory control under all conditions. Timing and method of application, weather and crop conditions, and other influencing factors are beyond the control of seller. Buyer assumes all risks of use, storage and handling of this product not in strict accordance with the directions and precautions given herewith.

Southern Agricultural Insecticides, Inc.

PALMETTO, FLA. HENDERSONVILLE, N. C. BOONE, N. C.
U. S. D. A. REG. NO. 829-145 B

NET WEIGHT _____

1362
NO. 1362
REGISTERED