

SA BRAND 50[®]

5.0% MANEB

5.0% SEVIN*

TOMATO and POTATO DUST

ACTIVE INGREDIENTS:

Maneb (manganese ethylenebisdithiocarbamate) 5.0%

Carbaryl (1-naphthyl N-methylcarbamate) 5.0%

INERT INGREDIENTS 90.0%

100.0%

*U. S. Pat. Nos. 2,903,478 - 2,504,404 - 1,710,822

Keep Out of Reach of Children

CAUTION! Harmful if swallowed or inhaled. May irritate eyes, nose, throat, and skin. Avoid breathing dust. Avoid contact with skin and eyes. Wash thoroughly after handling. Do not contaminate feed or foodstuffs. Store in a safe dry place. Burn empty container. Stay out of smoke and fumes.

May cause injury to tender plant parts especially when application is uneven or when foliage is damp. Control of the indicated insects and diseases may not be satisfactory during periods of frequent and heavy rainfall, when disease or insect infestations are extremely severe or when applications are uneven, inadequate or poorly timed.

Toxic to fish. Do not contaminate any body of water nor allow drift from the crop area treated. Apply this product only as specified on this label. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with the directions and precautions given herewith.

Southern Agricultural Insecticides, Inc.

PALMETTO, FLA. BOONE, N. C. HENDERSONVILLE, N. C.

EPA Reg. No. 829-147

E

NET WEIGHT

FOR USE ON

TOMATOES for the control of early and late blights, gray leaf spot (*Stemphyllium*), tomato fruit worms, flea beetles, tomato hornworms, and stink bugs. Apply 30 to 40 lbs. per acre at weekly intervals and following rains, beginning when plants are established in the field. For the control of late blight on maturing tomatoes apply 50 lbs. per acre when applied by ground duster and 60 lbs. per acre when applied by aircraft. Do not use within 5 days of harvest.

POTATOES for the control of early and late blights, fleabeetles, Colorado potato beetles, and leafhoppers. Make 30 to 40 lbs. per acre applications beginning when plants are about two inches high. Repeat at weekly intervals and following rains. For aircraft dust applications to control late blight on maturing potatoes apply 60 lbs. of SA-50 Brand Manzate Dust per acre.

BEANS for the control of Mexican bean beetles, Japanese beetle adults, corn earworms, and leafhoppers, and as an aid in the control of bean rust. Apply 20 to 40 lbs. per acre beginning when true leaves appear and repeat at weekly intervals. Do not use on beans within four days of harvest.

CABBAGE (in cabbage plant beds) for the control of downey mildew and fleabeetles. Dust thoroughly using 40 lbs per acre applications beginning when plants have begun to germinate and repeat at three-day intervals until plants are set in field. Do not use on cabbage within seven days of harvest.

PEPPER for the control of anthracnose, *Cercospora* leaf spot, corn earworms, fleabeetles, and hornworms. Apply 20 to 40 lbs. per acre. Begin applications when plants have become established in field, and repeat at weekly intervals and following rains.

ROSE, GLADIOLUS, and ZINNIAS for the control of black spot on Roses, *Botrytis* on Gladiolus, *Alternaria* leaf spot on Zinnias, Japanese beetle adults, thrips, fleabeetles, and leafhoppers. Thoroughly dust at weekly intervals and following rains. For Roses begin applications when first leaf buds begin to unfold. When dusting apply also to ground around roses to help control black spot infection from old leaves. For Gladiolus begin applications when first leaves appear. Continue applications after cutting to reduce *Botrytis* infection of bulbs. For Zinnias begin applications when plants are established.

For protection of honey bees avoid unnecessary use during periods when honey bees are visiting crops. When necessary to use during such periods, warn beekeepers well in advance of such applications.

NOTE TO PHYSICIANS: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal.

EPA REG. NO. 829-147

E

ACCEPTED