

829-147

5.0% MANEB

5.0% SEVIN*

TOMATO and POTATO DUST

131784

ACTIVE INGREDIENTS

Maneb (manganese ethylenebisdithiocarbamate)
Carbaryl (1-naphthyl N-methylcarbamate)**

INERT INGREDIENTS

TOTAL

*Trade-mark of Union Carbide Corporation for 1-Naphthyl N-methylcarbamate
**U.S. Pat. No. 2,903,478

829-147
5.0%
90.0%
100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician immediately. Do not induce vomiting unless other treatment is not available. May cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air. Apply respiration if indicated.

IF ON SKIN: Wash promptly with soap and water. Rinse thoroughly.

IF IN EYES: Flush eyes thoroughly with water for at least 15 minutes. Call a physician immediately.

NOTE TO PHYSICIAN: Atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

EPA EST. NO. 829 FL 1 (3) 829 NC 1 (4) 829 NC 2 (5) Number following Est. No. corresponds to first digit in Lot No. on bottom of bag.

Southern Agricultural Insecticides, Inc.

Palmetto, FL

Hendersonville, NC

Boone, NC

EPA REG. NO. 829-147

AA

NET WEIGHT 5 LBS

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

TOMATOES for the control of early and late blights, gray leaf spot (*Stemphylium*), tomato fruit worms, flea beetles, tomato hornworms, and stink bugs. Apply 30 to 40 lbs. per acre at weekly intervals and following rains, beginning when plants are established in the field. For the control of late blight on maturing tomatoes apply 60 lbs. per acre when applied by ground duster and 60 lbs. per acre when applied by aircraft. Do not use within 5 days of harvest.

POTATOES for the control of early and late blights, flea beetles, Colorado potato beetles, and leafhoppers. Make 30 to 40 lbs. per acre applications beginning when plants are about two inches high. Repeat at weekly intervals and following rains. For aircraft dust applications to control late blight on maturing potatoes apply 60 lbs. of SA 50 Brand Marzate Dust per acre.

BEANS for the control of Mexican bean beetles, Japanese beetle adults, corn earworms, and leafhoppers, and as an aid in the control of bean rust. Apply 20 to 40 lbs. per acre beginning when true leaves appear and repeat at weekly intervals. Do not use on beans within four days of harvest.

CABBAGE (in cabbage plant beds) for the control of downy mildew and flea beetles. Dust thoroughly using 40 lbs. per acre applications beginning when plants have begun to germinate and repeat at three-day intervals until plants are set in field. Do not use on cabbage within seven days of harvest.

PEPPER for the control of anthracnose, *Cercospora* leaf spot, corn earworms, flea beetles, and hornworms. Apply 20 to 40 lbs. per acre. Begin applications when plants have become established in field, and repeat at weekly intervals and following rains.

ROSE, GLADIOLUS, and ZINNIAS for the control of black spot on Roses, *Botrytis* on Gladiolus, *Alternaria* leaf spot on Zinnias, Japanese beetle adults, thrips, flea beetles, and leafhoppers. Thoroughly dust at weekly intervals and following rains. For Roses begin applications when first leaf buds begin to unfold. When dusting apply also to ground around roses to help control black spot infection from old leaves. For Gladiolus begin applications when first leaves appear. Continue applications after cutting to reduce *Botrytis* infection of bulbs. For Zinnias begin applications when plants are established.

NOTE: Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s).

Pesticide should be stored in the original container and placed in a locked storage area. Dispose of pesticide waste by securely wrapping the original container in several layers of newspaper and discard in trash. Do not reuse bag. Discard bag in trash.

May cause injury to tender plant parts especially when application is uneven or when foliage is damp. Control of indicated insects and diseases may not be satisfactory during periods of frequent and heavy rainfall, when disease or insect infestations are extremely severe or when applications are uneven, inadequate or poorly timed.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with the directions and precautions given herewith.

EPA REG. NO. 829-147

AA