

DIRECTIONS—Apply only when soil temperatures at 6" depth are 40° to 80° F. Soil should be in good tilth, with trash removed and clods broken. Immediately after applying, cover furrows and seal chisel channels and covered furrows by pressure rolling or flooding. Apply preplanting treatments 2 weeks before planting or setting. Do not disturb treated soil for 2 weeks after treatment. Do not feed treated plant parts to livestock. Do not use on plants inside homes.

Commercial growers should consult their state extension nematologist for specific application procedures consistent with this label.

TREATMENTS FOR INDICATED PLANTS — PRE-PLANTING-OVERALL—Apply in furrows or chisel channels 12" apart and 6" to 8" deep. Use 5 gal. per acre—1.46 oz. per 100 lineal feet of channel or 3 tablespoonsful diluted with 2 gallons of water per 100 lineal feet of furrow.

PRE-PLANTING — ROW TREATMENT—Apply in single chisel channel or furrow in center of row using same dosages per 100 lineal feet as indicated above.


POST PLANTING — DRENCH — FOR ESTABLISHED WOOD ORNAMENTALS AND TURF—Aerate soil with fork, rod or mechanical aerator. Apply 5 gallons per acre using 3 tablespoonsful in 19 gallons of water to cover 100 sq. ft. Sprinkle area with clear water after drenching. Keep children and pets off treated areas until Nemagon has been washed into soil.

ORNAMENTALS—Use dosage rates as indicated above. African Violet, Begonia, Caladium, Croton, Dieffenbachia, Dracaena, Haworthia, Ivy, Palm (Neantha Belle), Tradescantia, Roses — Pre Planting treatment as indicated — Post Planting Treatment only after roses have been established a year.

(Continued on right panel)



KEEP OUT OF REACH OF CHILDREN
WARNING: Harmful if inhaled. May burn skin. Causes irritation of eyes, nose and throat. May be absorbed through skin. Do not breathe vapor. Do not take internally. Use in a well ventilated area. Do not get in eyes. Avoid contact with skin or clothing. Wash thoroughly with soap and water after handling and before eating or smoking. In case of spillage on person or clothing, immediately remove contaminated shoes and clothing and wash with soap and water with plenty of water. Wash clothing. Wash clothes allowed call a physician. 2 tablespoonsful of this material has been washed into soil. Do not contaminate foodstuffs. Keep away from flame. Do not contaminate clothing. Do not use on this material.



DIRECTIONS—Apply only when soil temperatures at 6" depth are 40° to 80° F. Soil should be in good tilth, with trash removed and clods broken. Immediately after applying, cover furrows and seal chisel channels and covered furrows by pressure rolling or flooding. Apply preplanting treatments 2 weeks before planting or setting. Do not disturb treated soil for 2 weeks after treatment. Do not feed treated plant parts to livestock. Do not use on plants inside homes.

Commercial growers should consult their state extension nematologist for specific application procedures consistent with this label.


TREATMENTS FOR INDICATED PLANTS — PRE-PLANTING-OVERALL—Apply in furrows or chisel channels 12" apart and 6" to 8" deep. Use 5 gal. per acre—1.46 oz. per 100 lineal feet of channel or 3 tablespoonsful diluted with 2 gallons of water per 100 lineal feet of furrow.

PRE-PLANTING — ROW TREATMENT—Apply in single chisel channel or furrow in center of row using same dosages per 100 lineal feet as indicated above.

POST PLANTING — DRENCH — FOR ESTABLISHED WOOD ORNAMENTALS AND TURF—Aerate soil with fork, rod or mechanical aerator. Apply 5 gallons per acre using 3 tablespoonsful in 19 gallons of water to cover 100 sq. ft. Sprinkle area with clear water after drenching. Keep children and pets off treated areas until Nemagon has been washed into soil.

ORNAMENTALS—Use dosage rates as indicated above. African Violet, Begonia, Caladium, Croton, Dieffenbachia, Dracaena, Haworthia, Ivy, Palm (Neantha Belle), Tradescantia, Roses — Pre Planting treatment as indicated — Post Planting Treatment only after roses have been established a year.

(Continued on right panel)



(Continued from left panel)

FOR THE FOLLOWING USE 1 1/2 TIMES the dosage rates indicated. Arbor Vitae, Bleeding Heart (Dicentra), Boxwood, Camellia, Chrysanthemum (Hardy), Delphinium, Forsythia, Gardenia, Hibiscus, Illex (Holly), Juniper, Ligustrum (Privet), Lilac, Peony, Philodendron, Pittosporum, Pothos, Shasta Daisy.

VEGETABLES—Preplanting Treatments only. Use 1/2 Dosage rates indicated under treatments.

Snapbeans, cucumbers, summer squash, carrots, radishes, turnips, cabbage, cauliflower, broccoli and okra. Transplanted tomatoes use broadcast treatment at 30% of indicated rate.

TURF—Use Broadcast Drench Treatment—Bermuda, St. Augustine, Bahia, and Centipede grasses. Apply at 2 times rate indicated for drench treatment. (1 quart in 200 gallons of water per 1,000 sq. ft.) Repeat application in 4 months.

For Bent Grasses use 1 1/2 times dosage rate indicated for drench treatment. (1 1/2 pints in 200 gal. of water per 1,000 sq. ft.) Repeat when necessary.

Destroy empty container and bury with waste Nemagon in an isolated area.

This product may not give satisfactory control under all conditions. Timing, method of application, weather conditions, re-contamination of the treated area, and other factors influencing the results obtained from the use of this material are beyond the control of the seller. Buyer assumes all risks of use, storage, or handling not in strict accordance with the directions and precautions given herewith.


HENDERSONVILLE, N. C. 28739
P. O. BOX 429
TELEPHONE 704/692-2233

BOONE, N. C. 28607
P.O. BOX 85
TELEPHONE 704/264-8843

SA-50® BRAND

PALMETTO, FLORIDA 33561
P. O. BOX 218
TELEPHONE 813-722-3485

Southern Agricultural Insecticides, Inc.



Southern Agricultural Insecticides, Inc.

HENDERSONVILLE, N. C. 28739
P. O. BOX 429
TELEPHONE 704 692-2233

BOONE, N. C. 28607
P. O. BOX 85
TELEPHONE 704 264-8843

SA-50 BRAND

A C C O M P A N Y

SOUTHERN AGRICULTURAL INSECTICIDES, INC., HENDERSONVILLE, N. C.

S A BRAND 5 0 0

Nemagon® Emulsible Concentrate

FOR THE CONTROL OF
SOIL INFESTING NEMATODES
(Contains 2 quarts or 8.6
ACTIVE INGREDIENTS PER GALLON)
AND CERTAIN OTHER
N. MINERAL SOILS
Soil Fungicidal Per Gallon

66.0%
31.4%
31.8%

and water; for eyes flush with clean water. Do not use on plants inside homes. Do not use on plants inside homes. Do not use on plants inside homes.

Southern Agricultural Insecticides, Inc.
Palmetto, Florida
U.S.D.A. Reg. No. 829-156

WARNING: Keep out of reach of children. Do not use on plants inside homes. Do not use on plants inside homes. Do not use on plants inside homes.

Insecticides, Inc.
Hendersonville, N. C.
Contents: Liquid

DIRECTIONS: Apply only when soil temperatures at a depth of 10 to 20 inches should be in good till with fresh removed and chisel plow. Immediately after applying, cover furrows and soil chisel channels and covered furrows by pressure rolling or flooding. Apply on planting treatments 2 weeks before planting or sowing. Do not disturb treated and 2 weeks after treatment. Do not feed treated gentler to livestock. Do not use on plants inside homes.

TREATMENTS FOR INDIVIDUAL PLANTS:
PRE-PLANTING OVERSEED: Apply in furrows, chisel channels, or open and close chisel channels at 2 to 3 gallons per 100 lineal feet of furrow. Apply 2 to 3 gallons per 100 lineal feet of furrow.

PRE-PLANTING ROW TREATMENT: Apply in chisel channels or furrows in rows at 2 to 3 gallons per 100 lineal feet of furrow.

POST-PLANTING BROADCAST: For established wood, ornamentals, and turf. Apply with tank, not an mechanical sprayer. Apply 2 to 3 gallons per 100 lineal feet of furrow. Apply with clear water after drainage. Keep children and pets off treated areas until Nemagon has been washed into soil.

ORNAMENTALS: Use dosage rates as indicated above. Apply to: Arbor Vitae, Begonia, Caladium, Cyclamen, Daffodils, Freesia, Gladiolus, Hyacinth, Iris, Jasmine, Lilies, Tulips, and other plants. Do not use on plants inside homes.

S A BRAND 5 0 0

Nemagon® Emulsible Concentrate

FOR THE CONTROL OF ROOT KNOT AND CERTAIN OTHER
SOIL INFESTING NEMATODES ON MINERAL SOILS
(Contains 2 quarts or 8.6 lbs. Nemagon Soil Fungicidal Per Gallon)

66.0%
31.4%
31.8%

KEEP OUT OF REACH OF CHILDREN

WARNING: Harmful if inhaled. May burn skin. Causes irritation of eyes, nose and throat. May be absorbed through skin. Do not breathe vapor. Do not take internally. Use in a well ventilated area. Do not get in eyes. Avoid contact with skin or clothing. Wash thoroughly with soap and water after handling and before eating or drinking. In case of spillage on person or clothing, immediately remove contaminated shoes and clothing and wash with soap and water; for open flesh with plenty of water for at least 15 minutes and get medical attention. Wash clothing and do not wear until thoroughly cleaned. Do not use on plants inside homes. Do not use on plants inside homes. Do not use on plants inside homes.

Southern Agricultural Insecticides, Inc.
Palmetto, Florida
U.S.D.A. Reg. No. 829-156

(Continued from left panel)

FOR THE FOLLOWING USE 1. LIME: Use dosage rates indicated. Apply to: Arbor Vitae, Begonia, Heart (Quercus), Boxwood, Camellia, Citrus, Gardenia, Hibiscus, Holly, Juniper, Ligustrum (Privet), Lily, Peony, Philodendron, Poinsettia, Rhododendron, Shasta, Tuberose.

VEGETABLES: Pre-planting treatments and use. Dosage rates indicated in label treatments.

SNAPPELS: cucumber, summer squash, eggplant, radish, turnips, radishes, potatoes, broccoli and other transplanted and direct seeded treatment at 2 to 3 gallons per 100 lineal feet of furrow.

HERB: Use dosage rates as indicated above. Apply to: Bermuda, St. Augustine, Bahia, and other grasses. Apply at 2 to 3 gallons per 100 lineal feet of furrow. Do not use on plants inside homes.

FOR BENT GRASS: Use dosage rates as indicated above. Apply to: Bermuda, St. Augustine, Bahia, and other grasses. Apply at 2 to 3 gallons per 100 lineal feet of furrow. Do not use on plants inside homes.

Destroy empty container and label with water. Do not reuse.

This product may not be used on plants inside homes. Do not use on plants inside homes. Do not use on plants inside homes.