

**SA** BRAND **50**

# CAT AND DOG FLEA AND TICK DUST

**Also for Control of Horn Flies on Horses**

**ACTIVE INGREDIENTS:**

|  |                |
|--|----------------|
| Carbaryl (1-Naphthyl N-methyl-carbamate) * | 5.00%          |
| <b>INERT INGREDIENTS</b>                   | <b>95.00%</b>  |
|  | <hr/>          |
|  | <b>100.00%</b> |

\* U.S. Patent Nos. 2,903,478; 3,009,855.

**KEEP OUT OF REACH OF CHILDREN.**

**CAUTION!** Harmful if inhaled, swallowed, or absorbed through the skin. Avoid breathing dust. Avoid skin contact. Wash thoroughly after using and before eating or smoking.

**NOTE TO PHYSICIAN:** Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal.

DO NOT apply to animals or treat in or around animal quarters when animals are under medication, or when animals or animal quarters are under treatment from another cholinesterase inhibiting compound. Animals under treatment should be examined occasionally for skin irritation or other adverse symptoms. If reaction to treatment occurs, suspend treatment and consult a veterinarian.

BUYER ASSUMES all risks and liability from use, handling, or storage of this product not in strict accordance with the directions and precautions given herewith.

**Southern Agricultural Insecticides, Inc.**

Palmetto, Fla. Boone, N. C. Hendersonville, N. C.

U.S.D.A. Reg. No. 829-159 3 NET WEIGHT

To control fleas and ticks on cats and dogs. Dust liberally over entire animal, avoiding the eyes, and rubbing thoroughly to skin. Comb out dead ticks a few hours after treatment. Repeat at weekly intervals if needed. Do not treat nursing animals, small puppies, kittens under 4 weeks of age, or animals in a weakened condition.

In and around dog and cat quarters to control ticks and fleas — Dust in and around sleeping quarters, kennels and dog houses by means of shaker can or duster. Force dust into cracks and crevices. Repeat at weekly intervals as necessary to maintain control. When treating animals quarters do not contaminate feed, water, or utensils. When treating animal quarters, inside dwellings, apply in such a manner that small children cannot come in contact with dust. Do not treat bedding or quarters occupied by nursing animals, animals in a weakened condition, kittens under 4 weeks old, or young puppies.

**Horn flies on Horses:** Rub 2 ounces per animal into hair on back and neck. Repeat as necessary, but not more often than once every 4 days.

U.S.D.A. Reg. No. 829-159 8