

S A BRAND **50**[®]

Wettable or Dusting Sulfur

ACTIVE INGREDIENTS:

Sulfur 90.0%

INERT INGREDIENTS

10.0%

100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION: May cause irritation of eyes and skin. When dusting keep fire and sparks away from duster and dust stream.

PRECAUTION IN USING

Sulfur may cause injury to open blooms. Do not combine sulfur with oil sprays. Application of sulfur should not be closer than 3 weeks to an oil application and this period should be extended during dry or cold weather. Applications during very hot weather 90° F. or above may cause injury. Before applying commercial growers should consult proper U.S.D.A. or State Agricultural Authority for specific application procedures and dosages. Do not apply more than 50 lbs. per acre to Beans, 125 lbs. per acre to Citrus or 170 lbs. per acre to Peaches in any one application.

SUGGESTED FOR USE ON

Beans as an aid in the control of red rust. Apply at 4-7 day intervals using 40 lbs. per acre as a dust, or spray to obtain thorough coverage using 16 lbs. per 100 gallons (1/2 cup full per gallon). Begin applications when rust first appears and continue thereafter through harvest.

Aster, Azalea, Calendula, Chrysanthemum (hardy), Grape-myrtle, Dogwood, Dahlia, Delphinium, Ligustrum, Lilac, Maple, Phlox, Rhododendron, Rose, Snapdragon and Zinnia to control powdery mildew. Apply to obtain thorough coverage as a dust, or spray thoroughly using 4 lbs. per 100 gallons (2 tablespoonfuls per gallon). Make applications at 4 to 7 day intervals beginning when powdery mildew appears and continue applications as long as disease threatens.

Peaches to control brown rot. Apply to obtain thorough coverage using 5 to 6 lbs. per 100 gallons (3 tablespoonfuls per gallon). Apply at 3 day intervals during blossom period. Apply again as soon as all petals have fallen and repeat when 75% of the shucks have come off. Repeat applications thereafter at 10 to 14 days intervals to harvest. Applications may be continued through harvest period if brown rot is present or if humid or rainy weather occurs.

Citrus as a supplemental control of rust mite. Apply to obtain thorough coverage. Using 5 lbs. per 100 gallons (3 tablespoonfuls per gallon) as a spray, or as dust using 1/2 to 1 1/2 lbs. per tree depending on size of trees.

NON-WARRANTY Seller makes no warranty, expressed or implied as to the use of this material. Buyer assumes all risks and liability for results obtained by use or handling of this material whether or not in accordance with the directions or claimed so to be.

Southern Agricultural Insecticides, Inc.

Palmetto, Fla. Boone, N. C. Hendersonville, N. C.

U.S.D.A. Reg. No. 829-163

Net Weight 5 lbs.

REGISTERED

MAR 1971