



50

# WETTABLE OR DUSTING SULFUR

**ACTIVE INGREDIENTS:**

Sulfur.....	90.0%
INERT INGREDIENTS .....	10.0%
TOTAL.....	100.0%

**KEEP OUT OF REACH OF CHILDREN****CAUTION****STATEMENT OF PRACTICAL TREATMENT**

IF SWALLOWED: Contact a physician

IF INHALED: Remove victim to fresh air. Contact physician if irritation persists.

IF IN EYES: Flush eyes for 15 minutes with plenty of water, while lifting upper and lower lids. Seek medical attention if irritation persists

IF ON SKIN: Remove contaminated clothing and wash all exposed skin surfaces with soap and water. Contact a physician if irritation develops.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

May be harmful if swallowed. Avoid inhalation of dust. Avoid contact with skin and eyes. May cause allergic response. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from treated areas.

**PHYSICAL OR CHEMICAL HAZARDS**

When dusting, keep fire and sparks away from duster and dust stream.

EPA EST. NO. 829-FL-1 (3); 829-NC-1 (4); 829-NC-2 (5) Number following Est. No. Corresponds to first digit in Lot No. on bottom of bag

**Southern Agricultural Insecticides, Inc.**

Palmetto, FL 33581    Hendersonville, NC 28793    Boone, NC 28607  
EPA Reg. No. 829-163    CC    Net Weight: 2 lbs.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**REENTRY STATEMENT:**

Do not enter treated area within 24 hours after application of this product. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. (Indicate specific oral warnings which inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure.) When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: **CAUTION: area treated with Sulfur on (date of application). "Do not enter without appropriate protective clothing for 24 hours."** In case of accidental exposure wash with plenty of soap and water, flush eyes with water for at least 15 minutes, get medical attention.

**BEANS** as an aid in the control of rust. Apply at 4-7 day intervals using 40 lbs per acre as a dust, or spray to obtain thorough coverage using 16 lbs per 100 gallons (½ cup full per gallon). Begin applications when rust first appears and continue thereafter through harvest.

**ASTER, AZALEA, CALENDULA, CHRYSANTHEMUM (hardy), CRAPE-MYRTLE, DOGWOOD, DAHLIA, DELPHINIUM, LIGUSTRUM, LILAC, MAPLE, PHLOX, RHODODENDRON, ROSE, SNAPDRAGON AND ZINNIA** to control powdery mildew. Apply to obtain thorough coverage as a dust or spray thoroughly using 4 lbs per 100 gallons (2 tablespoonfuls per gallon). Make applications at 4 to 7 day intervals beginning when powdery mildew appears and continue applications as long as disease threatens.

**PEACHES** to control brown rot. Apply to obtain thorough coverage using 5 to 6 lbs per 100 gallons (3 tablespoonfuls per gallon). Apply at 3 day intervals during blossom period. Apply again as soon as all petals have fallen, and repeat when 75% of the shucks have come off. Repeat applications thereafter at 10 to 14 day intervals to harvest. Use lower rate when disease presence is light. Use higher rate during periods of high rainfall and heavy disease pressure. Applications may be continued through harvest period if brown rot is present or if humid or rainy weather occurs.

**CITRUS** as a supplemental control of rust mite. Apply to obtain thorough coverage using 5 lbs per 100 gallons (3 tablespoonfuls per gallon) as a spray, or as dust using ½ to 1½ lbs per tree depending on size of trees.

**NOTE:** Sulfur may cause injury to open blooms. Do not combine sulfur with oil sprays. Applications of sulfur should not be closer than 3 weeks to an oil application and this period should be extended during dry or cold weather. Applications during very hot weather 90°F or above may cause injury. Before applying this product consult your U.S.D.A. or State Agricultural Authority for specific application procedures and dosages. Do not apply more than 60 lbs per acre to Beans, 125 lbs per acre to Citrus or 170 lbs per acre to Peaches in any one application.

**PROCESSING NOTE:** When growing crops for processing, consult processor before applying sulfur.

**STORAGE AND DISPOSAL**

Store only in original container in a cool dry place inaccessible to children and pets and away from seed and foodstuffs. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash.

**CONTAINER DISPOSAL:** Do not reuse empty bag. Discard bag in trash.

**NOTICE:** Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS FOR USE and PRECAUTIONARY STATEMENTS stated on this label.

EPA REG. NO. 829-163

BB

**ACCEPTED****JAN 5 1987**

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 829-163