

GENERAL INFORMATION

To be applied by uniform broadcasting for the pre-emergence control of certain weed grasses including smooth and hairy crabgrass, foxtail, and barnyard grass.

For use on the following grasses in newly seeded or established turf: Blue grass, fescue, redtop, smooth brome, perennial ryegrass, orchardgrass, and the following strains of bentgrass: Penncross, seaside, highland, astoria, nimisila, C-1, C-7, and C-19.

Do not use on other bentgrass strains nor on bermuda grass, as injury to turf may result. The effect of SA-50 Brand Tupersan Crabgrass Killer on golf course greens may vary with turf management practices and environmental conditions; therefore, its use on golf greens is not recommended, as injury to the turf may result. Treatment will not control annual bluegrass (*poa annua*) clover, or most broadleaf weeds.

SA-50 Brand Tupersan Crabgrass Killer may be applied at any stage of turf development, including new seeding. At least 1/2 inch of water must be provided by rainfall or irrigation within 3 days after treatment in order to carry the chemical into the root zone of germinating weeds.

SA BRAND 50[®] TUPERSAN^{*} CRABGRASS KILLER GRANULES

*Trade Mark of E. I. DuPont de Nemours and Co., Inc., for: Siduron (1-(2-Methylcyclohexyl)-3-phenylurea). U. S. Patent Pending

ACTIVE INGREDIENTS:

Siduron [1-(2-Methylcyclohexyl)-3-phenylurea]	1.5%
INERT INGREDIENTS	98.5%
	100.0%

Keep Out of Reach of Children

CAUTION! May irritate eyes, nose, throat, and skin. Avoid breathing dusts. Avoid contact with skin, eyes, and clothing. After handling wash thoroughly and change to clean clothing.

Note: Use only on turf areas. Do not apply to crops used for food or feed purposes.

FOLLOW DIRECTIONS CAREFULLY. Timing and use of application, weather and lawn conditions, and other influencing factors in the use of this product are beyond control of the seller. Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with the directions given herewith.

SOUTHERN AGRICULTURAL INSECTICIDES, INC.

HENDERSONVILLE, N. C. BOONE, N. C. PALMETTO, FLA.
U. S. D. A. REG. NO. 829-178

ACCEPTED
JUL 7 1967
UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 829-178

NEW SPRING SEEDINGS

Apply as a final operation following seeding. Use 25 lbs. of SA-50 Brand Tupersan Crabgrass Killer per 3000 square feet.

NEW FALL SEEDINGS AND ESTABLISHED TURF

Apply in the spring just before expected emergence of annual weed grasses. Use 25 lbs. of SA-50 Brand Tupersan Crabgrass Killer per 2000 square feet.

All spreaders should be calibrated before using. To calibrate determine the spread width (swath) of the spreader. Place one lb. (1 quart) of granules in the spreader and apply granules in a straight line. Determine the distance covered. Adjust spreader setting and repeat the process until the spreader will deliver one lb. over distance given on the chart for desired dosage and spread width.

Feet of Travel by Spreader at Dosage

Spread Width	of 25 lbs. per 3000 Sq. Ft.	of 25 lbs. per 2000 Sq. Ft.
5 ft.	24	16
4 ft.	30	20
3 ft.	40	26
2 ft.	60	40

NET WEIGHT 25 LBS.

PM 25

829-178

7/7/67

73181