

CAUTION: Do not stain feed, foodstuffs, water, dishes, milk, milking equipment, or feed, feed and water utensils. Do not allow children or pets near treated surface until surface has dried.

HOUSEHOLD INSECTS—Apply as a 1/2% spray, prepared by mixing 2 1/2 fluid oz. of Vapona concentrate per gallon of water. Paint or apply as a coarse wet spray to limited areas only. Do not apply to entire walls or floor. If food or dish shelves are treated, allow to dry, then cover with shelf paper before replacing articles.

ANTS, COCKROACHES, AND SILVERFISH—Paint or spray baseboards, cabinets, closets, door and window casings and other hiding places including places where pipes go through walls and floors.

TICKS (including brown dog ticks)—Treat as above, also treat edges of carpets and behind pictures and draperies. Supplement by treating dogs with a product labeled for such use.

SPIDERS—Spray webs and corners frequented by spiders.

BED BUGS—Apply to tufts and seams only on mattress. Also apply to slats, springs, and bed frames. Dry at least 48 hours before use.

WASPS—Spray nest directly, preferably after dark without lights. After treatment of underground nests cover openings with moist dirt.

TO AID IN CONTROL OF HOUSE FLIES, STABLE FLIES, AND FACE FLIES

CATTLE—BEEF AND DAIRY—DIRECT APPLICATION—Apply as a 1% spray prepared by mixing 1 gallon of Vapona Concentrate in 25 gal. of water (5 oz. per gal. of water). Apply 1 to 2 fluid ounces per animal as a fine mist spray daily with a hand or automatic sprayer. Spray to thoroughly cover all parts of the animal body including back, flanks, and legs. Do not wet skin. Do not apply in excess of 2 fluid ounces per animal per day.

AGRICULTURAL PREMISES—DAIRY BARN—FEEDLOTS, MILK ROOMS, COUNTRY HOUSES AND STABLES.

INSIDE BUILDINGS Apply a 1% solution as a mist at the rate of one pint per 1,000 cu. ft. Do not apply to areas where animals have received direct application.

TO ALL INTERIOR AND EXTERIOR SURFACES Apply a 1/2% solution, prepared by mixing 1 gal. of Vapona Concentrate in 50 gal. of water (2 1/2 oz. per gal. of water), at rate of 1 quart per 1,000 sq. ft.

TO MANURE PILES AND GARBAGE DUMPS apply a 1/2% solution at rate of 2 quarts per 100 sq. ft.

GREENHOUSE—ORNAMENTAL PLANTS—As an aid to control mites and aphids when greenhouse is being heated by hot water or steam pipes. Make a fine mist to hot water or steam pipes by dribbling Concentrate on pipes at rate of 2 fluid ounces per 10,000 cu. ft. of greenhouse space. This treatment may be phytotoxic to some plant species, especially when flowers are showing color. Open Shasta and Fiji chrysanthemums are particularly sensitive.

BUYER assumes all risks of use, storage, or handling of this material not in strict accordance with the directions and precautions given herewith.

S A BRAND 50®
***VAPONA® EMULSIBLE CONCENTRATE**

(Contains 2 lbs. Dichlorovos (D.D.V.P.) per gallon)

ACTIVE INGREDIENTS	
(2,2-Dichlorovinyl dimethyl phosphate)	21.9%
Related Compounds	1.5%
Heavy aromatic naphtha solvent	65.0%
INERT INGREDIENTS	11.6%
	100.0%

This product is sold on the condition that it will be used only by properly equipped personnel who are familiar with the use and dangers of Dichlorovos (D.D.V.P.). This product is intended for use by licensed pest control operators, industrial and agricultural users—it is not intended for use by home owners. See side panels for uses and additional safety precautions.

DANGER! POISON!

KEEP OUT OF REACH OF CHILDREN



POISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. REPEATED INHALATION OR SKIN CONTACT, MAY, WITHOUT SYMPTOMS, PROGRESSIVELY INCREASE SUSCEPTIBILITY TO VAPONA INSECTICIDE POISONING. Do not swallow or get in eyes, on skin, or on clothing. Do not breathe spray mist or vapors. Wear clean natural rubber gloves; clean, water-proof protective clothing and goggles. Replace gloves frequently and destroy used gloves. Wear a (D.D.V.P.) protection. Wash thoroughly with soap and water after handling and before eating or smoking. If material gets into eyes, immediately flush with water for at least 10 minutes and get medical attention. In case of contact, remove all contaminated clothing and wash skin with soap and water; wash clothing before reuse. Decontaminate areas and equipment by washing with strong lye solution, then destroy used containers as indicated to prevent re-use.

® Trademark, Shell Chemical Corp.

SOUTHERN AGRICULTURAL INSECTICIDES, Inc.

PALMETTO, FLORIDA

BOONE, N. C.

HENDERSONVILLE, N. C.

U.S.A. Reg. No. 879-179

NET LIQUID CONTENTS 1 GALLON

ACCEPTED
 OCT 16 1967
 UNDER FEDERAL INSECTICIDE
 FUNGICIDE AND RODENT CONTROL ACT
 FOR ECONOMIC POISON REG. DIST.
 ED UNDER NO. 429-179

RED
INK

RED
INK

