

SA 50

KILL


RATS AND MICE

ACTIVE INGREDIENTS:

3-(Alpha-Acetylfurfuryl)-4-Hydroxycoumarin 0.025%

INERT INGREDIENTS 99.975%

100.000%



See Back Panel for Additional Precautions and Directions

KEEP OUT OF REACH OF CHILDREN

CAUTION! If swallowed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of Vitamin K combined with blood transfusions may be indicated, as in the case of hemorrhage caused by overdose of 4-hydroxycoumarin.

Southern Agricultural Insecticides, Inc.

PALMETTO, FLA. BOGALUS, LA. C HENDERSONVILLE, N. C.
NEW WT.

DIRECTIONS

This bait contains an anti-coagulant chemical. Its remarkable killing power depends on its being consumed regularly in small doses for a period of days, usually 5 to 14.

Do not contaminate feed or foodstuffs. Store in a safe place. Protect bait stations from children, pets, and domestic animals by using bait boxes or suitable covers. Care should be exercised, when placing bait stations in locations near stored feed or foodstuffs, to avoid contamination.

Place bait in a protected spot where the rats and mice ordinarily feed and water, or where they travel or live. Baits should be placed along walls and in corners rather than in the middle of the room. Use shallow containers for the bait.

For RATS use one or two bait placements (1/2 to 1 pound each) at every point of large activity — look for such tell-tale signs as gnawings, tracks or trails, burrows, droppings, or damage.

For MICE a large number of bait placements should be made. Locate tablespoonful quantities of bait at 6 to 10 ft. intervals where mice are active.

Keep Feeding Stations Well Supplied

Inspect and replenish each station daily for the first week and regularly thereafter until bait is no longer being taken. Never allow a feeding station to remain empty for over 24 hours. If some bait stations are untouched, move to a better location. If baits become sour or moldy (as might occur during extended periods of hot, humid weather), replace with fresh bait as needed.

Where re-infestation from dumps and fields may occur, permanent bait stations should be set out and replenished as needed.

Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with the directions given herewith.