



BAIT PELLETS KILL Land Snails and Slugs


— FOR USE ON —

Ornamentals, Certain Vegetables and Tobacco

ACTIVE INGREDIENTS

Metaldohyde	3.25%
Carbaryl (1-Naphthyl N-methylcarbamate)	4.25%
INERT INGREDIENTS	92.50%
	100.00%

Contains Sevin — Reg. T. M. Union Carbide Corporation for 1-Naphthyl N-methylcarbamate. U.S. Pat. No. 2,903,478



KEEP OUT OF REACH OF CHILDREN

WARNING! May be fatal if swallowed. May cause irritation of the skin. Avoid contact with skin, eyes and clothing. Avoid breathing dust. Wash thoroughly after using. Keep container closed. Store in a safe, cool, dry place away from heat and open flame. Dispose of empty container by burning. Stay out of smoke and fumes.

For Commercial Use Only

This pesticide may be fatal to children and dogs or other pets if eaten. Keep children and pets out of treated area.
EPA EST 829 FL 1-31-829 NC 1-4-829 N.C. 5-1-829
Responsible for distribution in N.C.

Southern Agricultural Insecticides, Inc.

PALMETTO, FLA. BOONE, N. C. HENDERSONVILLE, N. C.
EPA Reg. No. 829-182 C

DIRECTIONS

For use in the Southeastern United States as an aid in the control of land snails and slugs.

Apply to soil surface around plants. In green houses where tomatoes and ornamentals are grown. In plant beds, fields, and gardens where tobacco, ornamentals, ornamental turfs, and the following food crops are grown: Beans, broccoli, brussel sprouts, cabbage, cauliflower, eggplant, lettuce, okra, pepper, and tomatoes.

Distribute pellets evenly around plants using 1 lb. per 1000 square feet. When applying to areas planted in food crops, do not contaminate edible parts of plants. Do not apply to lettuce within 14 days of harvest, to broccoli, brussel sprouts, cabbage or cauliflower within 3 days of harvest.

When slugs are moving from a known area into a planting, pellets may be placed in a "guard line" at the edge of the planting. Pellets should be applied in late afternoon. Pellets will be less effective when placed in very dry locations. Applications should be repeated at 10-day intervals as necessary to maintain control. Three applications at 10-day intervals should cover a complete life cycle of snails or slugs.

Keep children, pets and domestic animals from treated areas until pellets have disintegrated.

This product is toxic to wildlife. Birds and other wildlife in treated areas may be harmed. Apply this product only as specified on this label. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with the directions given herewith.

