



BAIT PELLETS

KILL LAND SNAILS AND SLUGS

FOR USE ON ORNAMENTALS, CERTAIN VEGETABLES AND TOBACCO



ACTIVE INGREDIENTS:	
Metaaldehyde (2,4,6,8-tetramethyl-1,3,5,7-tetraoxycyclo-octane)	3.25%
Carbaryl (1-Naphthyl N-methylcarbamate)	4.25%
INERT INGREDIENTS	92.50%
TOTAL	100.00%

KEEP OUT OF REACH OF CHILDREN
CAUTION

FIRST AID N-METHYL CARBAMATE

IF SWALLOWED: Call a physician immediately. Gastric lavage is indicated if material was taken internally. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

IF IN EYES: Flush eyes with plenty of water. Call a physician immediately.

IF ON SKIN: Wash immediately with soap and water. Rinse thoroughly.

IF INHALED: Remove victim to fresh air. Apply respiration if indicated.

NOTE TO PHYSICIAN: Atropine sulfate is antidotal.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be fatal if swallowed. May cause irritation of the skin. Avoid contact with skin, eyes, and clothing. Avoid breathing dust. Keep container closed. This pesticide may be fatal to dogs or other pets if eaten. Keep pets out of treated area.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants and shoes plus socks. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should: wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, pets, birds, and other wildlife. For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

Store in a safe, cool, dry place away from heat and open flame.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store this pesticide in original container in a locked storage area.

PESTICIDE DISPOSAL (HOUSEHOLD ONLY): Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL (HOUSEHOLD ONLY): Do not reuse container. Discard empty bag in trash.

PESTICIDE DISPOSAL (EXCEPT HOUSEHOLD): Wastes resulting from the use of this product may be disposed of on site or at an approved waste facility.

CONTAINER DISPOSAL (EXCEPT HOUSEHOLD): Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned stay out of smoke.

Southern Agricultural Insecticides, Inc.

Palmetto, Fla. 34220 Hendersonville, N.C. 28793 Boone, N.C. 28607

EPA REG. No. 829-182 J3 **Net Weight: 2.5 lbs. (1.13 kg)**

PM 04 829-182
7/22/99 M182

ACCEPTED
JUL 22 1999
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 829-182

NEW LABEL

282

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is coveralls, waterproof gloves, and shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Keep children and pets off treated area until subsequent rainfall or irrigation has washed the bait into the soil and the area is dried.

For use as an aid in the control of land snails and slugs.

ORNAMENTALS AND TURF GRASSES

For use in greenhouses, plant beds, fields, and gardens on ornamentals, and ornamental turfs: Apply to soil surface around plants. Distribute pellets evenly around plants using 1 lb. per 1,000 square feet. When slugs are moving from a known area into a planting, pellets may be placed in a "guard line" at the edge of the planting. Pellets should be applied in late afternoon. Pellets will be less effective when placed in very dry locations. Applications should be repeated at 10-day intervals as necessary to maintain control. Three applications at 10-day intervals should cover a complete life cycle of snails or slugs.

CROPS RECOMENDATIONS

Apply to soil surface around plants. In greenhouses where tomatoes and ornamentals are grown. For use in greenhouses on Tomatoes¹ and in plant beds, fields, and gardens on: Beans¹, Broccoli¹, Brussels Sprouts¹, Cabbage¹, Cauliflower¹, Eggplant¹, Lettuce¹, Okra¹, Pepper¹, Tobacco¹, and Tomatoes¹. Distribute pellets evenly around plants using 1 lb. per 1,000 square feet. When applying to areas planted in food crops, do not contaminate edible parts of plants. Do not apply to lettuce within 14 days of harvest, or to broccoli, brussels sprouts, cabbage or cauliflower within 3 days of harvest. When slugs are moving from a known area into a planting, pellets may be placed in a "guard line" at the edge of the planting. Pellets should be applied in late afternoon. Pellets will be less effective when placed in very dry locations. Applications should be repeated at 10-day intervals as necessary to maintain control.

¹ Applications must not exceed 4 lbs. of A.I. (123 lbs. of bait) per Acre per growing season.

² Applications must not exceed 3.2 lbs. of A.I. (98 lbs. of bait) per Acre per growing season.

NOTE: Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s).

Follow instructions carefully. Timing and care of application, weather and lawn conditions, and other factors influencing the results obtained by the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with the directions and precautions given herewith.



0 51538 10910 9

FPA ES1 No. 829-FL-1(3); 879-NC-1(4); 829-NC-2(5) Number following establishment No. corresponds in first digit in lot number.