

For use in the Southeast to control European red, apple rust, Willamette, two spotted (red spider) strawberry (Atlantic), citrus red (purple), clover (brown almond), six-spotted, spruce, and tomato russet mites. Kelthane E.C. is compatible with most insecticides and fungicides that are compatible with solvent based formulations. It is not compatible with lime or other alkaline materials. Kelthane should be applied to thoroughly cover all above ground parts of the plant, especially the undersides of the leaves. To prepare spray thoroughly mix 1 quart of Kelthane in 100 gallons of water (2 teaspoonfuls per gallon), except where other dosages are indicated. Certain mites may show resistance to Kelthane treatments in your area and applications may not give satisfactory control under all conditions. Applications to open blooms, and to damp foliage may cause injury.

Apples, Crabapples, and Quinces. Make 2 or more applications as needed at 10 to 14 day intervals when mites appear. Apply preferably in the second or later cover sprays. Do not apply within 7 days of harvest. Do not exceed 2½ gal. per acre in any one application.

Peaches, Apricots, Nectarines, Cherries, Plums, Prunes. Apply at petal fall or later, when mite population appears. Repeat application in 30 days (if needed) (but not sooner). Do not exceed 2 gal. per acre in any one application. Do not apply to cherries, plums or prunes within 7 days of harvest or to peaches, apricots, or nectarines within 14 days of harvest.

Grapes, Raspberries, Dewberries, Blackberries, Boysenberries. Apply after bloom, when mites appear. Repeat in 7 to 10 days if necessary. Do not exceed ¾ gal. per acre in any one application. Do not apply to grapes within 7 days of harvest or to raspberries, dewberries, blackberries and boysenberries within 2 days of harvest.

S A BRAND 5 O® KELTHANE® E. C.

CONTROLS MANY SPECIES OF MITES
Contains 1.6 lbs. Kelthane per gallon)

ACTIVE INGREDIENTS

1,1-bis(chlorophenyl)-2,2,2-Trichloroethanol ... 18.5%
Xylene 13.0%
INERT INGREDIENTS 8.5%

U.S. Patent Nos. 2,812,362 and 2,812,280 100.0%

Kelthane® is a registered Trademark of Rohm & Haas Company.

KEEP OUT OF REACH OF CHILDREN

CAUTION: Avoid contact with skin, eyes, and clothing. Harmful if swallowed. Avoid inhalation of spray mist. Do not smoke while spraying. Wash thoroughly after handling. Do not contaminate feed or feedstuffs. Do not use inside the home. Rinse, destroy, and bury empty containers and spray waste in an isolated area away from water supplies. Inflammable — keep away from heat and open flame.

TOXIC TO FISH. KEEP OUT OF ANY BODY OF WATER.

Southern Agricultural Insecticides, Inc.

Hendersonville, N. C. Palmetto, Fla. Boone, N. C.

U.S.D.A. Reg. No. 829-184 Net Contents 16 oz. Liquid

Strawberries. Apply to foliage as needed for mite control, using 5 gallons of diluted spray per 1000 sq. ft. For cyclamen mite make 3 applications, at 10 day intervals, when new growth starts in spring. Apply with high pressure using 10 gallons of diluted spray per 1000 sq. ft. Do not apply within 2 days of harvest.

Citrus — for fall and winter combined control of spider mite and rust mites and for control of spider mites wherever control is necessary. Apply 2½ pints per 100 gallons (2½ tsp. per gal.) and repeat as necessary. Do not apply within 7 days of harvest. Do not exceed 5 gal. per acre in any one application.

Clover mites entering buildings. Spray, from the outside, to thoroughly cover windowsills, walls, foundations, plants at the base of infested buildings and adjacent lawn area. Repeat applications in 10 to 14 days if needed.

Ornamentals — Arboretum, azalea, begonia, Boston fern, boxwood, cactus, chrysanthemum (outdoor), coleus, crabtree, cyclamen, dahlia, day lily, dogwood, easter lily, elm, euonymus (European), evergreen fern, fuchsia, gardenia, geranium, gladiolus, hamamelis (Eastern), hydrangea, iris, ivy, juniper, laurel, lilac, magnolia, maple, oak, oleander, petunia, philodendron, redbud, rose, snapdragon, spruce, violet (African), willow and wisteria. Apply when mites appear and repeat if necessary.

NOTICE: Follow instructions carefully. Timing and application are important. Do not mix with other chemicals, and other influencing factors in the use of this product are beyond control of the seller. Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with the directions and precautions given herewith.

