

Suggestions for Use: Thorough coverage is essential. Spray to obtain thorough coverage of trunks, limbs and leaves. Do not apply to plants that are near wilting or wilting from lack of moisture. Do not apply with sulfur or lime-sulfur, or within 3 weeks before or after application of sulfur or lime-sulfur. Apply only on clear mild days, before late afternoon. Avoid spraying during periods of high temperature (above 90° F.). Do not exceed two applications per year. Do not apply in Florida after October 1st. Do not apply when temperatures fall below, or are expected to fall below 42° F. within 3 weeks of application. Except as otherwise indicated, additional applications, or applications of other spray oil formulations when necessary, should be at least 6 weeks apart. Avoid spraying open blooms.

The oil used in this formulation meets Florida Citrus Standards designated as F. C. 435-66.

Azalea, Boxwood, Camellia, Gardenia, Ligustrum, Pittosporum and Viburnum for control of camellia scale, ewonymus scale, Florida red scale, Florida wax scale and spider mites. Apply preferably when scale crawlers and mites first appear using 2 quarts per 100 gals. (4 teaspoonfuls per gal.). Repeat if necessary. Do not apply to easily injured sensitive plants as sugar and Japanese Maple, beech, hickory, walnut and butternut.

S A BRAND 5 O® ETHION-SUPERIOR OIL

CONTAINS .67 POUNDS ETHION PER GALLON

ACTIVE INGREDIENTS	
Ethion (0,0,0-Tetraethyl S,S-methylene bisphosphorodithioate)	9.0%
Petroleum Oil (Superior type)	75.0%
INERT INGREDIENTS	16.0%
U.S. Patent No. 2,873,228	100.0%

KEEP OUT OF REACH OF CHILDREN

WARNING: May be fatal if swallowed, inhaled or absorbed through skin. Do not breathe vapor or spray mist. Do not get in eyes or on skin or clothing. In case of skin contact, wash with soap and water. For eyes flush with plenty of water and get medical attention. Wash clothing with soap and hot water before re-use. Wash thoroughly after using and before eating or smoking. Do not contaminate food or feed products. Do not store at temperatures below 32° F. This product is poisonous to fish and wildlife. Do not contaminate any body of water. Do not apply when weather conditions favor drift from the area.

See side panels for directions and additional precautions.

Southern Agricultural Insecticides Inc.

Palmetto, Fla. Hendersonville, N. C. Boone, N. C.

Net Contents 16 oz. Liquid

Florida Citrus: For the control of citrus red mite, citrus rust mite, Texas citrus mite, black scale, chaff scale, Florida red scale, glover scale, purple scale, snow scale and yellow scale. Apply preferably in June or July, using 2 quarts per 100 gallons (4 teaspoonfuls per gal.). Do not apply to kumquats, citrus citron, or navel oranges. Do not apply more than once per season nor within 21 days of harvest to lemons or limes. Do not apply more than twice per crop to tangerines. Do not repeat applications to citrus within 90 days. Do not exceed 2,000 gallons of indicated spray mixture per acre in any one application.

Camellia: For tea scale. Apply when crawlers first appear using 3 quarts per 100 gal. (2 tablespoonfuls per gal.). Repeat application if necessary.

Junipers: For the control of two spotted mite apply when first detected. For juniper scale treat immediately after eggs hatch using 2 quarts per 100 gal. (4 teaspoonfuls per gal.).

Follow instructions carefully. Timing, method of application, weather conditions, mixtures with other chemicals and other influencing factors in the use of this product are beyond control of the seller. Buyer assumes all risks of use, storage, or handling not in strict accordance with the directions and precaution given herewith.