

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician immediately. Induce vomiting by giving one or two glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Get medical attention immediately. Remove patient from area of exposure. If breathing has stopped, give artificial respiration.

IF ON SKIN: Remove contaminated clothing. Wash affected area with soap and water. If person is unconscious or incoherent, transport to hospital immediately.

IF IN EYES: Flush eyes with water for 15 minutes. Seek medical attention.

NOTE TO PHYSICIAN: Atropine sulfate is antidotal. 2-PAM is also antidotal and should be administered in conjunction with atropine.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: May be fatal if swallowed, inhaled or absorbed through skin. Do not breathe vapor or spray mist. Avoid contact with skin. Wash thoroughly after handling and before eating or smoking. Avoid contamination of food or feed.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, wildlife and shrimp. Do not apply directly to water or wetlands. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

CITRUS & ORNAMENTAL SPRAY

Use 1/4 to 2 tablespoonfuls per gallon of water
Contains Ethion and Oil

ACTIVE INGREDIENTS:

Ethion (0,0,0-D⁺ - Tetraethyl S,S' - Methylene bisphosphorodithioate) 8.4%

Petroleum Oil (Superior type) 86.1%

INERT INGREDIENTS 5.5%

TOTAL 100.0%

CONTAINS 825 POUNDS ETHION PER GALLON.

The oil used in this formulation meets Florida Citrus Standards designated as F.C. 435-68.

**KEEP OUT OF REACH OF CHILDREN
WARNING**

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA EST NO 829-FL-1 (3), 829-NC-1 (4), 829-NC-2 (5) Number following Est. No. corresponds to first digit in Lot No. on side of label

Southern Agricultural Insecticides, Inc.

Palmetto, FL 34220 Hendersonville, NC 28733 Boone, NC 28607
EPA Reg No. 829-186 J1 Net Contents: 1 Pint

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Azalea, Berried, Camellia, Gardenia, Ligustrum, Philadelphus and Viburnum for control of camellia scale, acyrthosus scale, Florida red scale, Florida wax scale and spider mites. Apply preferably when scale crawlers and mites first appear using 1 1/2 tablespoonfuls per gallon of water. Repeat if necessary. Do not apply to easily injured sensitive plants as sugar and Japanese maple, beech, hickory, walnut and butternut.

Camellia: For tea scale. Apply when crawlers first appear using 2 tablespoonfuls per gallon of water. Repeat application if necessary.

Jambur: For the control of two spotted mite, apply when first detected. For russet scale, treat immediately after eggs hatch using 1 1/2 tablespoonfuls per gallon of water.

Florida Citrus: For the control of citrus red mite, citrus rust mite, Texas citrus mite, black scale, chaff scale, Florida red scale, glover scale, purple scale, snow scale and yellow scale. Apply preferably in June or July, using 1 1/2 tablespoonfuls per gallon of water. Do not apply to tangerines or citrus groves. Do not apply more than once per season nor within 21 days of harvest to lemons or limes. Do not apply more than twice per crop to tangerines. Do not repeat applications to citrus within 90 days.

Thorough coverage is essential. Spray to obtain thorough coverage of trunks, limbs and leaves. Do not apply to plants that are near wilting or withering from lack of moisture. Apply only on clear, mild days, before late afternoon. Avoid spraying during periods of high temperature. Do not exceed two applications per year. Do not apply in Florida after October 1st. Do not apply when temperatures fall below, or are expected to fall below 42°F. Except as otherwise indicated, additional applications, or applications of other spray or formulations when necessary, should be at least 6 weeks apart. Avoid spraying open blooms.

NOTE: Do not apply this product through any type of irrigation system.

STORAGE AND DISPOSAL

Pesticide Storage: Store pesticide in the original container in a locked storage area. Do not store below 32°F.

Pesticide Disposal: Securely wrap original container in several layers of newspaper and discard in trash.

Container Disposal: Do not reuse container. Rinse thoroughly before discarding in trash.

Buyer assumes all risks of use, storage or handling not in strict accordance with the directions and precautions given hereon.

ACCEPTED

FEB 03 1989

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, this pesticide is registered under EPA Reg No. 829-186