DIRECTIONS FOR USE

App'y only as a dilute spray when weather conditions favor good coverage. Do not apply when temperatures are below 40° F. or when freezing or near freezing temperatures or frost will occur within 24 hours. Do not use in combination with sulfur, lime sulfur, captan folpet or dich-Ione. Do not apply within 7 days of an application of captan, folpet or dichlone. Do not apply within 30 days after or 7 days before an application of sulfur or lime sulfur. Do not apply to plants that are under stress from lack of moisture. Add to spray tank when at least 3/4 full. Agitate thoroughly and continue agitation. Do not allow mixture to stand. Do not combine with other spray materials unless prior use of the mixture has proven compatible. If oil or oil in combination with other spray materials does not mix or emulsify properly, do not use as plant injury may result. Spray to obtain complete thorough plant coverage.

APPLES --- for early season control of European Red Mite and San Jose' Scale. Mix 2 gallons of oil per 100 gallons of water. Do not excede 25 gallons of oil per acre. For European Red Mite apply preferably as a delayed dormant spray. For San Jose Scale application may be made as a dormant or delayed dormant spray. NOTE: In North Carolina, apply when buds show 1/2-Inch new growth (second pre-bloom spray).

50® SA SUPERIOR 70-SECOND EMULSIFIABLE SPRAY OIL

ACTIVE INGREDIENTS: Mineral Oil INERT INGREDIENTS

(Unsulfonated Residue, ASTM-D-483 96%)

Keep Out of Reach of Children

CAUTION! Harmful if swallowed. If Spray Oil, with-out other pesticide additives is swallowed, do not induce vomiting. May cause skin irritation. When used in combination with other pesticides which may be absorbed through the skin, absorbtion of these pesticides may be enhanced. Wash thoroughly after using. Change to clean clothing daily.

This product is formulated with Orchex 796. A product of Humble Oil and Refining Company, Houston, Texas.

SEE SIDE PANELS FOR INSTRUCTIONS



NET CONTENTS LIQUID 55 GALS.

99.5% 00.5% 100.0%

ь. <u>з</u>

ັບ

71

DORMANT SPRAYS FOR ORNAMENTALS

Dormant sprays are generally more effective in the spring just before new growth starts.

PLANT		Quarts per 100 gal.	Table spoonf per ga	luls
Azulea	Bark Scale	6	4	On early blooming varieties, apply dur- ing warm weather in late fall.
Camellia	Camellia Leaf Sca	le 6	4	Supplement to sum- mer applications.
Holly	Holly Scale	8	6	In spring before buds open.
Pittosporum	Cottony Cushion Scale	8	6	2 applications.
Privet	White Peach Scale	e 12	8	2 applications 2 weeks apart.
Pyracantha	Wax Scale	12	8	Before visible growth starts.
Rose	Rose Scale	6	4	Before visible growth starts.

U.

ſ

BR

Þ

Ζ

 $\textcircled{\baselinetwidth}$

NOTE: This material may not give satisfactory control under all conditions. Thoroughness of application, weather conditions, severity of insect infestations, and other factors influencing the results obtained from the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling not in strict accordance with the directions and precautions given herewith.

U. S. D. A. Reg. No. 829-187