

ACCEPTED
 Apr 12 1971
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR REG. OR REG. PENDING REG. NO. 09,217, 345
 TO ATTACHED COMMENTS

A BRAND 5 0®
**25% LINDANE WETTABLE
 SPRAY CONCENTRATE**

For use by commercial growers only. Do not use in or around home.

ACTIVE INGREDIENTS	
Gamma isomer of benzene hexachloride from Lindane	25.0%
INERT INGREDIENTS	75.0%
	100.0%

FOR USE ON Cucumbers, Melons, Pumpkins and Squash, to control aphids. Apply to obtain thorough coverage using 1 lb. 25% Lindane per 100 gallons spray. Use 200 gallons per acre on maturing crops with normal growth, proportionately less on younger crops or crops with sparse vine growth. Do not exceed 2 lbs. 25% Lindane per acre. Do not apply Lindane more than one time per year. Additional treatments for aphids may be required. Consult your Cooperative Extension Service for an approved material. Remove excess residue by washing or brushing. Do not apply to fields to be planted to potatoes, peanuts, or other root crops within 2 years as Lindane may adversely affect their flavor. Late applications of Lindane may cause off flavor in melons.

Highly toxic to fish and wildlife. Keep out of any body of water. Do not apply when conditions favor drift from the treated area. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning equipment or disposal of waste. Apply only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Extension Service.

KEEP OUT OF REACH OF CHILDREN

WARNING: May be fatal if swallowed. Readily absorbed through skin. Do not breathe spray mist. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling, and change to clean clothing. In case of skin contact, wash with soap and water. For eyes, flush with water and get medical attention. Store in a safe place. Do not contaminate feed or foodstuffs. Destroy empty container. Bury with waste pesticide in an isolated area.

Do not exceed indicated dosages. Before applying, consult proper state extension authority to determine if local recommendations are consistent with this label. This product can cause plant injury and may not give satisfactory control under all conditions. Timing, method of application, mixtures with other chemicals, weather conditions, severity of insect infestation and reinfestation, and other factors influencing the results obtained from the use of this product are beyond control of the seller.

Buyer assumes all risks of use, storage, or handling of this product not in strict accordance with the directions and precautions given herewith.

Southern Agricultural Insecticides, Inc.

Hendersonville, N.C. Palmetto, Fla. Boone, N.C.

Net Weight _____

U.S.D.A. Reg. No. 829-191

Finished label out size: 4" x 6 1/8"