

ACCEPTED

1 MAR 1988

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 829-196

DIPEL® DUST

ACTIVE INGREDIENTS:

Bacillus thuringiensis Berlin, Var. Kurstaki potency of 200 International Units per milligram.....0.04%

INERT INGREDIENTS:.....99.96%

TOTAL 100.00%

*Equivalent to 90 Million International Units per pound.

© Trademark of Abbott Laboratories.

KEEP OUT OF REACH OF CHILDREN

CAUTION:

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May cause irritation of eyes, nose, throat, skin. Avoid inhalation: Wash thoroughly after handling.

ENVIRONMENTAL HAZARDS

Keep out of lakes, ponds or streams. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a cool dry area. Do not store for extended periods of time or where temperatures are above 90°F.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.

EPA EST. NO. 829-FL-1 (3); 829-NC-1 (4); 829-NC-2 (5) Number following Est. No. corresponds to first digit in Lot No. on bottom of bag.

Southern Agricultural Insecticides, Inc.

Palmetto, FL 33581

Hendersonville, NC 28793

Boone, NC 28607

EPA Reg. No. 829-196

AA

Net Weight: 2 lbs.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Apply recommended amounts when air is calm, using ground equipment, or hand duster. Apply at first sign of infestation and repeat at weekly intervals as needed to maintain control.

To be effective Dipel Dust must be eaten by the insect. Thorough leaf coverage, especially to the underside of the leaves is essential. Some insects stop feeding after application and appear to be uneffected for several days after application. Check foliage for signs of fresh feeding if insects appear to remain active after applying.

Apply higher indicated dosages to maturing plants when applied by ground equipment. Do not exceed maximum dosage, nor use less than the lower dosage indicated.

Crop	Pest	Lbs. Per Acre
BROCCOLI	Cabbage looper	20-60
	Imported cabbageworm	16-50
CABBAGE	Cabbage looper	20-60
	Imported cabbageworm	16-50
CAULIFLOWER	Cabbage looper	20-60
	Imported cabbageworm	16-50
CELERY	Cabbage looper	20-60
COLLARDS	Cabbage looper	20-60
	Imported cabbageworm	16-50
CUCUMBERS	Cabbage looper	16-50
GRAPES	Grape leaf folder	16-50
Start treating as soon as possible after hatching and before larvae are protected by leaf folds. Apply by ground equipment using 16-50 pounds per acre to obtain thorough coverage of leaf surfaces.		
KALE	Cabbage looper	20-60
	Imported cabbageworm	16-50
LETTUCE	Cabbage looper	20-60
MELONS	Cabbage looper	16-50
MUSTARD GREENS	Cabbage looper	20-60
	Imported cabbageworm	16-50
POTATOES	Cabbage looper	20-60
TOBACCO	Tobacco hornworm	16-50
TOMATOES	Cabbage looper	20-60
	Tomato hornworm	16-50
TURNIP GREENS	Cabbage looper	20-60
	Imported cabbageworm	16-50

Dipel Dust is exempt from tolerance requirements and may be applied in the recommended amounts on indicated crops to day of harvest.

The manufacturer makes no warranties, expressed or implied, concerning this product or the results obtained from its use which extend beyond the description and precautions given herewith.

EPA Reg. No. 829-196

AA

829-196

63/01/1988

1/1