



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OCT 24 1995

OFFICE OF  
PREVENTION, PESTICIDES  
AND TOXIC SUBSTANCES

Mr. John C. Diem  
Southern Agricultural Insecticides  
P.O. Box 218  
Palmetto, FL 34220

Label Amendment Resubmission of dated 10/16/95 in  
Response to PR Notice 93-7  
EPA Reg. No. 829-196  
Thuricide HP 90M Dust Microbial Insecticide

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments reflected on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling
- AND
- WITHIN one year from date of this acceptance.


Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)  
Office of Pesticide Programs (7505C)  
U.S. Environmental Protection Agency  
401 M Street, SW  
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)  
Office of Pesticide Programs  
Room 266A, Crystal Mall 2  
1921 Jefferson Davis Highway  
Arlington, VA 22202

Sincerely,



Jim Tompkins, Deputy Chief  
Fungicide/Herbicide Branch  
Registration Division (7505W)

Attachment



50<sup>®</sup>

Home & Garden

# THURICIDE HP-90M\* DUST MICROBIAL INSECTICIDE

*# foliar*

24 1995

### ACTIVE INGREDIENTS:

Bacillus thuringiensis Berliner, Var. Kurstaki potency of 200 International Units per milligram*	0.04%
<b>INERT INGREDIENTS:</b>	99.96%
<b>TOTAL</b>	100.00%

\*Equivalent to 90 Million International Units per pound.

## KEEP OUT OF REACH OF CHILDREN CAUTION

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** May cause irritation of eyes, nose, throat, and skin. Avoid inhalation. Wash thoroughly after handling.

#### ENVIRONMENTAL HAZARDS

Keep out of lakes, streams, and ponds. For terrestrial uses, do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

### STORAGE AND DISPOSAL

**PESTICIDE STORAGE:** Store in a cool dry area. Do not store for extended periods of time or where temperatures are above 90°F.

**PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash.

**CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

EPA EST. NO.829-FL-1 (3); 829-NC-1 (4); 829-NC-2 (5) *Reg # 827 196*  
Number following Est. No. corresponds to first digit in Lot No. on bottom of bag.

**Southern Agricultural Insecticides, Inc.**

Palmetto, FL 34220 Hendersonville, NC 28793 Boone, NC 28607

## DIRECTIONS FOR USE

It is a violation Federal Law to use this product in a manner inconsistent with its labeling. Apply recommended amounts when air is calm using a hand duster. Apply at first sign of infestation and repeat at weekly intervals as needed to maintain control. To be effective Dipel Dust must be eaten by the insect. Thorough leaf coverage, especially to the underside of the leaves is essential. Some insects stop feeding after application and appear to be unaffected for several days after application. Check foliage for signs of fresh feeding if insects appear to remain active after applying. Apply higher dosage to maturing plants. Do not exceed maximum dosage, nor use less than the lower dosage.

CROP	PEST	OZS. PER 1,500 SQ. FT.
BROCCOLI	Cabbage Looper	11 - 33
	Imported Cabbageworm	9 - 27 1/2
CABBAGE	Cabbage Looper	11 - 33
	Imported Cabbageworm	9 - 27 1/2
CAULIFLOWER	Cabbage Looper	11 - 33
	Imported Cabbageworm	9 - 27 1/2
CELERY	Cabbage Looper	11 - 33
	Imported Cabbageworm	9 - 27 1/2
COLLARDS	Cabbage Looper	11 - 33
	Imported Cabbageworm	9 - 27 1/2
CUCUMBERS	Cabbage Looper	9 - 27 1/2
GRAPES	Grape Leaf Folder	9 - 27 1/2
<b>NOTE:</b> Start treating as soon as possible after hatching and before larvae are protected by leaf folds. Apply to obtain thorough coverage of leaf surfaces.		
KALE	Cabbage Looper	11 - 33
	Imported Cabbageworm	9 - 27 1/2
LETTUCE	Cabbage Looper	11 - 33
MELONS	Cabbage Looper	9 - 27 1/2
MUSTARD GREENS	Cabbage Looper	11 - 33
	Imported Cabbageworm	9 - 27 1/2
POTATOES	Cabbage Looper	11 - 33
TOMATOES	Cabbage Looper	11 - 33
	Tomato Hornworm	9 - 27 1/2
TURNIP GREENS	Cabbage Looper	11 - 33
	Imported Cabbageworm	9 - 27 1/2

Dipel Dust may be applied at the recommended rate up to the day of Harvest.

The manufacturer makes no warranties, expressed or implied, concerning this product or the results obtained from its use which extend beyond the description and precautions given herewith.

