


829-196

10/16/2000

1/2



Home & Garden®

# THURICIDE HP-90 DUST MICROBIAL INSECTICIDE

**ACCEPTED**

OCT 16 2000

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 829-196

## CONTAINS DIPEL® BIOLOGICAL INSECTICIDE

### ACTIVE INGREDIENTS:

Bacillus thuringiensis Subsp. Kurstaki solids, spores, and insecticidal toxins. ....	0.04%
INERT INGREDIENTS: .....	99.96%
TOTAL .....	100.00%

\*The percent active ingredient does not indicate product performance and potency. DIPEL® is a Trademark of VALENT

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if inhaled. Avoid breathing spray dust. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

#### ENVIRONMENTAL HAZARDS

Keep out of lakes, streams, and ponds. For terrestrial uses, do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

### FIRST AID

**If on skin:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes.  
**If in eyes:** Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center for treatment advice.  
**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center for treatment advice.

### STORAGE AND DISPOSAL

**PESTICIDE STORAGE:** Store in a cool dry area. Do not store for extended periods of time or where temperatures are above 90°F.  
**PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash.  
**CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

## Southern Agricultural Insecticides, Inc.

Palmetto, FL 34220      Hendersonville, NC 28793      Boone, NC 28607  
EPA REG. No. 829-196      J3      **Net Weight: 4 lbs. (1.81 kg)**

## DIRECTIONS FOR USE

It is a violation Federal Law to use this product in a manner inconsistent with its labeling. Apply recommended amounts when air is calm using a hand duster. Apply at first sign of infestation and repeat at weekly intervals as needed to maintain control. To be effective Dipel Dust must be eaten by the insect. Thorough leaf coverage, especially to the underside of the leaves is essential. Some insects stop feeding after application and appear to be unaffected for several days after application. Check foliage for signs of fresh feeding if insects appear to remain active after applying. Apply higher dosage to maturing plants. Do not exceed maximum dosage, nor use less than the lower dosage.


<b>CROP</b>	<b>PEST</b>	<b>OZS. PER 1,500 SQ.FT.</b>
<b>BROCCOLI</b>	Cabbage Looper	11 - 33
	Imported Cabbageworm	9 - 27 1/2
<b>CABBAGE</b>	Cabbage Looper	11 - 33
	Imported Cabbageworm	9 - 27 1/2
<b>CAULIFLOWER</b>	Cabbage Looper	11 - 33
	Imported Cabbageworm	9 - 27 1/2
<b>CELERY</b>	Cabbage Looper	11 - 33
	Cabbage Looper	11 - 33
<b>COLLARDS</b>	Cabbage Looper	11 - 33
	Imported Cabbageworm	9 - 27 1/2
<b>CUCUMBERS</b>	Cabbage Looper	9 - 27 1/2
	Cabbage Looper	9 - 27 1/2
<b>GRAPES</b>	Grape Leaf Folder	9 - 27 1/2
	Cabbage Looper	11 - 33
<b>KALE</b>	Cabbage Looper	11 - 33
	Imported Cabbageworm	9 - 27 1/2
<b>LETTUCE</b>	Cabbage Looper	11 - 33
	Cabbage Looper	9 - 27 1/2
<b>MELONS</b>	Cabbage Looper	9 - 27 1/2
	Cabbage Looper	11 - 33
<b>MUSTARD GREENS</b>	Cabbage Looper	11 - 33
	Imported Cabbageworm	9 - 27 1/2
<b>POTATOES</b>	Cabbage Looper	11 - 33
	Cabbage Looper	11 - 33
<b>TOMATOES</b>	Cabbage Looper	11 - 33
	Tomato Hornworm	9 - 27 1/2
<b>TURNIP GREENS</b>	Cabbage Looper	11 - 33
	Imported Cabbageworm	9 - 27 1/2

**NOTE:** Start treating as soon as possible after hatching and before larvae are protected by leaf folds. Apply to obtain thorough coverage of leaf surfaces.

Dipel Dust may be applied at the recommended rate up to the day of Harvest.

The manufacturer makes no warranties, expressed or implied, concerning this product or the results obtained from its use which extend beyond the description and precautions given herewith.

For information or questions contact our website at [Southernag.com](http://Southernag.com) or email us at [info@southernag.com](mailto:info@southernag.com)



0 51538 03674 0

EPA EST. NO. 829-FL-1 (3); 829-NC-1 (4); 829-NC-2 (5)

Number following Est. No. corresponds to first digit in Lot No. on bottom of bag.

EPA REG. No. 829-198 J3