

6-14-72
829-200

SA BRAND 50 SEVIN 10% DUST

ACTIVE INGREDIENTS:
Carbaryl (1-Naphthyl N-methylcarbamate) 10.0%
INERT INGREDIENTS 90.0%
100.0%

*Trademark of Union Carbide Corp. for:
1-Naphthyl N-methylcarbamate U. S. Patent No. 2,903,475

Keep Out of Reach of Children

CAUTION! Harmful if swallowed or inhaled. May irritate eyes, nose, throat and skin. Do not breathe dust. Avoid contact with skin, eyes, and clothing. Wear clean long sleeved work clothing daily. Wash thoroughly after using. Do not contaminate feed or foodstuffs. Burn empty container. Stay out of smoke and fumes.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Do not contaminate any body of water nor allow drift from the treated area. This product is highly toxic to bees exposed to direct treatment or residues on plants. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not exceed indicated dosages. Before applying consult proper state extension authority to determine if local recommendations are consistent with this label. This product may injure tender plant parts particularly when applications are uneven, or when plants are wet, or when applications are made during periods of sustained high humidity. This product may not give satisfactory control under all conditions. Timing, method of application, weather conditions, severity of insect infestation, and other factors influencing the results obtained from the use of this product are beyond control of seller.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with the directions and precautions given herewith.

Southern Agricultural Insecticides, Inc.

Hendersonville, N. C. Palmetto, Florida Boone, N. C.

U.S.A. Reg. No. 829-200

NET WEIGHT LBS:

On ornamentals and trees, apply dust to upper and lower leaf surfaces, and to the soil around the base of the plant. Repeat treatment as necessary.

Keep children away from treated areas until irrigation has washed dust into soil.

Woody ornamentals and ornamental yard trees, to control bagworm, birch leaf miner, blister beetles, Japanese beetle, and June beetle adults, mimosa webworm, orange tortix, periodical cicada, tent caterpillar, thorn bug, and exposed thrips: Apply dust lightly but thoroughly as needed to maintain control to upper and lower leaf surfaces.

Consult State Extension Specialist for timing of application.

Do not use on Boston ivy.

Corn: Fall armyworm and corn earworm (budworm damage only) apply 15 to 20 lbs. per acre, with ground equipment, directing the dust into the whorl. For insects attacking the silks and ears apply dust at a rate of 15 to 20 lbs. per acre, starting when silks first appear and repeat at 2 to 3-day intervals until silks begin to dry. Do not apply within 7 days of harvest for forage.

Application of this product to the tassel region of corn during pollen shed period will seriously reduce bee population.

For Grasses and Pasture: For the control of armyworms and thrips use 12 to 15 lbs. per acre.