



SEVIN®

BRAND CARBARYL INSECTICIDE

10% DUST

FOR USE ON CORN, WOODY ORNAMENTALS, AROUND THE OUTSIDE OF HOMES, HAY, PASTURES, AND LAWNS FOR CONTROL OF ARMYWORMS, CENTIPEDES, MILLIPEDES, EARWIGS, CERTAIN BEETLES, AND SOD WEBWORMS. ALSO FOR THE CONTROL OF FIRE ANT MOUNDS.

ACTIVE INGREDIENTS:

Carbaryl (1-Naphthyl N-methylcarbamate)	10.0%
INERT INGREDIENTS	90.0%
TOTAL	100.0%

SEVIN® is the registered trademark of Rhône-Poulenc for Carbaryl Insecticide.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT N-METHYLCARBAMATE

IF SWALLOWED: Call a physician immediately. Induce vomiting immediately. To induce vomiting give one or two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air. Apply respiration if indicated.

IF ON SKIN: Wash promptly with soap and water. Rinse thoroughly.

IF IN EYES: Flush eyes thoroughly with water for at least 15 minutes. Call a physician immediately.

NOTE TO PHYSICIAN: Atropine sulfate is antidotal.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants and shoes plus socks. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic and estuarine invertebrates. For terrestrial use, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

EPA EST NO. 029 FL-1 (3); 029 NC-1 (4); 029 NC-2 (5) Number following Est. No. corresponds to first digit in Lot No. on bottom of bag.

Southern Agricultural Insecticides, Inc.

Palmetto, FL 34220
EPA REG. NO. 029-200

Hendersonville, NC 28793
J2

Beane, NC 28007
Net Weight: 4 lbs.

ACCEPTED
MAR 20 1995

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 029-200

1482

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, Waterproof gloves, Shoes plus Socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Keep children and pets off treated areas until dust has settled.

NOTE: Where dosage ranges are indicated, use higher dosages on maturing plants and proportionately less when applied by ground equipment to smaller plants. Do not exceed maximum indicated dosages. Before applying consult proper State Extension Authority to determine if local recommendations are consistent with this label. This product may injure tender plant parts particularly when applications are uneven, or when plants are wet, or when applications are made during periods of sustained high humidity. This product may not give satisfactory control under all conditions. Timing, method of application, weather conditions, severity of insect infestation and other factors influencing the results obtained from the use of this product are beyond the control of the seller.

Yards and Outside Areas Around the home to control centipedes, millipedes and earwigs. Apply to grass, shrubbery, mulch, and foundations around the outside of the house. Use 2 lbs. of dust per 1000 sq. ft. Repeat only if necessary.

Ornamental Turfs to control sod webworms and fall armyworms: Treat as soon as worms are detected using 2 lbs. per 1000 sq. ft. Repeat treatment as necessary.

Keep children away from treated turf and border areas until rain or irrigation has washed dust into soil.

Imported Fire Ants—Mound Treatment: Lightly disturb the top of the mound in order to make the ants more active. Then immediately apply 1-2 ounces to the mound and the area surrounding the mound. For best results apply in the early morning or late afternoon. Applications made under prolonged hot and dry conditions may be ineffective due to the location of the ants deep within the nest. Do not severely disturb the mound as mound relocation may occur. Do not irrigate after treatment. Treat new mounds as they appear.

Woody Ornamentals and Ornamental Yard Trees, to control bagworm, birch leaf miner, blister beetles, Japanese beetle, and June beetle adults, mimosa webworm, orange tortix, periodical cicada, tent caterpillar, thorn bug, and exposed thrips: Apply dust lightly but thoroughly as needed to maintain control to upper and lower leaf surfaces.

Consult State Extension Specialist for timing of application.

Do not use on Boston Ivy.

Corn: Fall armyworm and corn earworm (budworm damage only) apply 15 to 20 lbs. per acre, with ground equipment, directing the dust into the whorl. For insects attacking the silks and ears apply dust at rate of 15 to 20 lbs. per acre, starting when silks first appear and repeat at 2 to 3-day intervals until silks begin to dry. Do not apply within 7 days of harvest for forage.

Application of this product to the tassel region of corn during pollen shed period will seriously reduce bee population.

For Grasses and Pasture: For the control of armyworms and thrips use 12 to 15 lbs. per acre.

NOTE: Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s).

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal. **STORAGE:** Store pesticide in the original container in a locked storage area. **PESTICIDE DISPOSAL:** To dispose of excess pesticide, securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with the directions and precautions given herewith.

EPA REG NO 829-200 J2



BEST COPY AVAILABLE