




CARBARYL 10% DUST

FOR USE ON CORN, WOODY ORNAMENTALS, AROUND THE OUTSIDE OF HOMES,
HAY, PASTURES, AND LAWNS FOR CONTROL OF ARMYWORMS, CENTIPEDES,
MILLIPEDES, EARWIGS, CERTAIN BEETLES, AND SOD WEBWORMS.
ALSO FOR THE CONTROL OF FIRE ANT MOUNDS.

ACTIVE INGREDIENT:	
Carbaryl (1-Naphthyl N-methylcarbamate)	10.0%
INERT INGREDIENTS	90.0%
TOTAL:	100.0%



**KEEP OUT OF REACH OF CHILDREN
CAUTION**

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or inhaled. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before use.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants and shoes plus socks. Follow manufacturers instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Note to reviewer: The following Residential Use Requirements section is for the residential use labeling:

RESIDENTIAL USE REQUIREMENTS

Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product. Change clothing as soon as possible after use. Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet. Do not allow children or pets to contact treated area until dust has settled.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water, or to areas where surface water is present or to Intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

EPA EST. NO. 829-FL-1 (3); 829-NC-1 (4); 829-NC-2 (5) Number following Est. No. corresponds to first digit in Lot No.

Southern Agricultural Insecticides, Inc.

Palmetto, FL 34220

Hendersonville, NC 28739

Boone, NC 28607

EPA REG. NO. 829-200 J5

Net Weight: 4 lb. (1.81 kg.)

FIRST AID N-METHYLCARBAMATE

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Never give anything by mouth to an unconscious person. **IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

NOTE TO PHYSICIAN: Carbaryl is an N-methyl carbamate insecticide which is a cholinesterase inhibitor. Specific antidote consists of atropine. Narcotics and other sedatives should not be used. Drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended.

Have product container or label with you when calling a poison control center or doctor or going for treatment.

For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378

Note to reviewer: The following WPS sections are for the commercial use labeling (over 25 Lbs.)

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls Waterproof gloves, Shoes plus Socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Keep children and pets off treated areas until dust has settled.

DIRECTIONS FOR USE

It is a violation of Federal law to use the product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not use this product on plants above chest height.

Use a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate duster as available. Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust. Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering. For plants above chest height, consider using a liquid carbaryl product. Do not repeat application more than once a week.

NOTE: Where dosage ranges are indicated, use higher dosages on maturing plants and proportionately less when applied by ground equipment to smaller plants. Do not exceed maximum indicated dosages. Before applying consult proper State Extension Authority to determine if local recommendations are consistent with this label. This product may injure tender plant parts particularly when applications are uneven or when plants are wet, or when applications are made during periods of sustained high humidity.

Note to reviewer: The following use rates are for the commercial use labeling.

Ornamental Turfs to control sod webworms and fall armyworms: Treat as soon as worms are detected using 2 lbs. per 1,000 sq. ft. Repeat treatment as necessary.

Imported Fire Ants - Mound Treatment: Lightly disturb the top of the mound in order to make the ants more active. Then immediately apply 1 - 2 ounces to the mound and the area surrounding the mound. For best results apply in the early morning or late afternoon. Applications made under prolonged hot and dry conditions may be ineffective due to the location of the ants deep within the nest. Do not severely disturb the mound as mound relocation may occur. Do not irrigate after treatment. Treat new mounds as they appear.

Woody Ornamentals to control bagworm, birch leaf miner, blister beetles Japanese beetle, and June beetle adults, mimosa webworm, orange tortix, periodical cicada tent caterpillar, thorn bug, and exposed thrips: Apply dust lightly but thoroughly as needed to maintain control to upper and lower leaf surfaces.

Consult State Extension Specialist for timing of application. Do not use on Boston Ivy or Trees.

Corn (sweet): Fall armyworm and corn earworm (budworm damage only) apply 15 to 20 lbs. per acre, with ground equipment, directing the dust into the whorl. For insects attacking the silks and ears apply dust at rate of 15 to 20 lbs. per acre, starting when silks first appear and repeat at 7 day intervals until silks begin to dry.

Application of this product to the tassel region of corn during pollen shed period will seriously reduce bee population. (Ears 2, forage 14, fodder 48)

For Grasses and Pasture: For the control of armyworms and thrips use 12 to 15 lbs. per acre. (14)

NOTE: Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s).

Number in () behind crop name is the days required between final application and harvest (PHI).

Note to reviewer: The following use rates are for the residential use labeling.

Yards and Outside Areas Around the home to control centipedes, millipedes and earwigs. Apply to grass, shrubbery, mulch, and foundations around the outside of the house. Use 2 lbs. of dust per 1,000 sq. ft. Repeat only if necessary.

Ornamental Turfs to control sod webworms and fall armyworms: Treat as soon as worms are detected using 2 lbs. per 1,000 sq. ft. Repeat treatment as necessary. Keep children away from treated turf and border areas until rain or irrigation has washed dust into soil.

Imported Fire Ants - Mound Treatment: Lightly disturb the top of the mound in order to make the ants more active. Then immediately apply 1 - 2 ounces to the mound and the area surrounding the mound. For best results apply in the early morning or late afternoon. Applications made under prolonged hot and dry conditions may be ineffective due to the location of the ants deep within the nest. Do not severely disturb the mound as mound relocation may occur. Do not irrigate after treatment. Treat new mounds as they appear.

Woody Ornamentals to control bagworm, birch leaf miner, blister beetles Japanese beetle, and June beetle adults, mimosa webworm, orange tortix, periodical cicada tent caterpillar, thorn bug, and exposed thrips: Apply dust lightly but thoroughly as needed to maintain control to upper and lower leaf surfaces. Consult State Extension Specialist for timing of application. Do not use on Boston Ivy or Trees.

Corn (sweet): Fall armyworm and corn earworm (budworm damage only) apply .4 to .5 lbs. per 1000 sq. ft., with ground equipment, directing the dust into the whorl. For insects attacking the silks and ears apply dust at rate of .4 to .5 lbs. per 1000 sq. ft., starting when silks first appear and repeat at 7 day intervals until silks begin to dry.

Application of this product to the tassel region of corn during pollen shed period will seriously reduce bee population. (Ears 2, forage 14, fodder 48)

For Grasses and Pasture: For the control of armyworms and thrips use .3 to .4 lbs. per 1000 sq. ft. (14)

NOTE: Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s).

Number in () behind crop name is the days required between final application and harvest (PHI).

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store pesticide in the original container in a locked storage area.

PESTICIDE DISPOSAL: To dispose of excess pesticide, securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.

Note to reviewer: The following Storage and Disposal is for the residential use labeling.

PESTICIDE STORAGE: Store in an area away from children and pets.

PESTICIDE DISPOSAL: If Empty: Do not reuse empty container. Put in trash or offer for recycling if available.

If partly filled: Call you local solid waste agency or 1-800-CLEANUP for disposal instructions Never place unused product down any indoor or outdoor drain.

WARRANTY AND LIMITATION OF DAMAGES

SELLER warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. BUYER assumes the risk of any use contrary to such directions. SELLER makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY, and no agent of SELLER is authorized to do so except in writing and with specific reference to this warranty. In no event shall SELLER'S liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

For information and questions concerning this product please contact us by email at support@southernag.com.