

BRAND

HORNFLY AND LICE DUST

USE ON CATTLE, SWINE AND GOATS

ACTIVE INGREDIENT:

Technical Methoxychlor 10%

(Equivalent to 8.8% 2,2-bis (p-methoxyphenyl)-
1,1,1-trichloroethane and 1.2% others isomers and
reaction products)

INERT INGREDIENTS 90%
100%

KEEP OUT OF REACH OF CHILDREN

CAUTION: Wash thoroughly after handling. Do not contaminate feed and water. Do not breathe dust.

TOXIC TO FISH: Keep out of lakes, ponds and streams.

Do not reuse bag. Bury when empty.

BEEF CATTLE (May be used on young dairy animals up to two weeks of freshening.) For hornfly, lice and fleas, thoroughly dust animal and work dust into hair. Use proportionally more or less for larger or smaller animals. Repeat application in 2-3 weeks, if needed.

SWINE for lice control, make liberal applications to animals and pens.

SHEEP AND GOATS for lice control make a thorough application to animal and work dust into hair. Applications to sheep, following shearing, gives temporary relief from sheep keds.

Do not apply to lactating dairy animals or goats.

DUST (BUTT) BAGS for hornfly control on beef cattle. Place 12 1/2 lbs of dust in each bag and use one bag for 10-15 animals. Dust bags may be commercially manufactured type or home made. For home made bags use double burlap sacks and cover burlap with polyethylene on the top and sides or hang in an area protected from rain. Suspend bags by heavy wire or chain in loafing areas and in water and feeding paths. Do not refill bags more often than every 4 weeks. Do not apply to animals being finished for slaughter. Do not use this treatment on dairy animals.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions and precautions given here-with.

Southern Agricultural Insecticides, Inc.

Palmetto, Fla.

Hendersonville, N.C.

Boone, N.C.

EPA Reg. No. 829-201

Net Weight