


[829-202]



ACCEPTED  
APR 22 1972  
UNDER FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO. 829-202

SHAKE WELL before using — Pour recommended amount into  $\frac{1}{2}$  filled spray tank. Agitate and maintain agitation. Do not allow dilute spray to stand in tank more than a few hours. Combinations with most fungicides and stickers are not deleterious if used promptly. A Non-ionic spreader sticker, used sparingly, is recommended when applications must be made when frequent rainfall and heavy dews are anticipated. Do not combine with alkaline materials such as bordeaux mixture or lime sulfur. During hot, sunny, or windy weather, late afternoon and early evening applications produce better results. Apply to obtain thorough leaf coverage. Thuricide must be carried into the insects gut while feeding, to be effective. Apply at first sign of infestation. Repeat application in five to seven days and thereafter at weekly intervals as necessary to maintain control.

Thuricide is exempt from tolerance on the indicated food crops at the recommended dosage and may be used to the day of harvest.

This package is intended for homeowner use. The active ingredient concentration is the same as Thuricide packaged for use by commercial growers, however commercial growers should follow label instructions on packages intended for commercial use.

**ORNAMENTALS.** For the control of insects listed in food crop section, use 4 teaspoonfuls per gallon.

pests = cabbage looper  
imported cabbage worm  
fruit tree leaf roller  
orange dog  
tobacco budworm  
tobacco hornworm

# S A BRAND 5 O<sup>®</sup> THURICIDE<sup>®</sup> - HPC

FOR THE CONTROL OF CERTAIN PLANT  
FEEDING CATERpillARS AND WORMS  
USE 1 to 4 TEASPOONFULS PER GALLON

**ACTIVE INGREDIENTS:**

*Barillus thuringiensis* Berliner, potency of  
4,000 International Units (at least 6  
million viable spores) per milligram\* 0.8%

Petroleum hydrocarbon solvent 3.0%

**INERT INGREDIENTS** 96.2%

\*Equivalent to 4.0 billion International Units per quart.

Thuricide is a registered trademark of International Minerals & Chemical Corp.

### KEEP OUT OF REACH OF CHILDREN

**CAUTION:** Do not take internally. Wash thoroughly after using.

Rinse container thoroughly with water and discard when empty.

**STORE IN A COOL PLACE:** Insect activity may be impaired by storage at temperatures above 90 degrees Fahrenheit. Protect from freezing.

Packaged and Distributed By

**Southern Agricultural Insecticides, Inc.**

Palmetto, Fla. Hendersonville, N.C. Boone, N.C.

EPA Reg. No. 829-202 Net Contents ..... Liquid

### RECOMMENDATIONS

THOROUGH COVERAGE IS ESSENTIAL. Apply indicated amount in each gallon of water and spray to obtain thorough coverage of foliage.

CROP	PEST	Teasp. Per Gal.
Beans (N. Carolina, Virginia)	Cabbage Looper	2
Broccoli	Cabbage Looper Imported Cabbageworm	4 3
Cabbage	Cabbage Looper Imported Cabbageworm	4 3
Cauliflower	Cabbage Looper Imported Cabbageworm	4 3
Collards, Kale	Cabbage Looper	3
Cucumbers (Southeast Only)	Cabbage Looper	2
Lettuce	Cabbage Looper	4
Melons	Cabbage Looper	4
Mustard Greens	Cabbage Looper	3
Oranges, Navel	Fruit Tree Leafroller	4
Oranges	Orange Dog	1
Potatoes	Cabbage Looper	4
Spinach (N. Carolina, Virginia)	Cabbage Looper	4
Tobacco	Cabbage Looper Tobacco Budworm Tobacco Hornworm	3 3 2
Tomatoes	Cabbage Looper Tomato Hornworm	2 2
Turnip Greens	Cabbage Looper	3
Watermelons (Florida)	Windworm Complex	2

