

S A BRAND 5 ®

Insecticide Bait

Contains 5% DYLOX®

ACTIVE INGREDIENT:

Dimethyl (2,2,2-trichloro-1-hydroxyethyl) phosphonate*

5%

INERT INGREDIENTS

95%

100%

U.S. Patent No. 2,701,225
 Dylox Reg. TM by Farbenfabriken
 Bayer A.G., Chemagro A Division of Baychem Corp., Licensee.

KEEP OUT OF REACH OF CHILDREN

CAUTION: May be harmful if swallowed, inhaled, or absorbed through the skin. Avoid contact with skin. Do not breathe dust. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse. Do not contaminate feed or food. If poisoning occurs, obtain prompt medical aid. Prolonged exposure will result in cholinesterase depression.

TO PHYSICIAN: Atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on label.

Do not use on other crops used for food or forage. Use only according to label directions. Applications at rates above those shown may result in illegal crop residue. Do not treat food crops grown in the greenhouse.

Rinse equipment and containers and dispose of wastes by burying in non-crop land away from water supplies. Dispose of empty container by burying with wastes or by burning. (Keep out of smoke).

Follow directions carefully. Timing and method of application, weather conditions and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with the directions and precautions given herewith.

Southern Agricultural Insecticides, Inc.

Palmetto, Fla. Hendersonville, N.C. Boone, N.C.

Net Weight

EPA Reg. No. 829-203 A

DIRECTIONS

5% Dylox Bait gives excellent control of surface feeding cutworms, salt marsh caterpillars, armyworms and crickets infesting many crops and ornamentals. Applications may be made as a band over planted row or around stems of plants in the row. If such treatments are not practical, apply as a broadcast treatment over the infested area. If the field has a history of previous infestations or if cutworms are suspected, apply 5% Dylox Bait immediately after planting or transplanting.

RECOMMENDED APPLICATIONS

	Pounds		REMARKS
	5% DYLOX BAIT Per Acre	Per 1,000 Sq. Ft.	
Beans (Dry and Snap)	30	3/4	Do not apply within 14 days of harvest. Do not feed treated vines to meat or dairy animals.
Beans (Lima)	30	1/4	Do not apply more than twice per season nor harvest for food, feed or forage within 30 days of treatment.
Corn	20	1/2	Apply once per season when plants are 3-12 inches high. Do not use corn for food, feed or ensilage within 40 days of treatment.
Cowpeas (Southern peas, blackeye peas, crowder peas)	30	3/4	Do not apply more than twice per season nor within 14 days of harvest. Do not feed treated vines to meat or dairy animals.
Brussel Sprouts Cabbage Cauliflower	20	1/2	Do not apply within 21 days of harvest.
Collards Lettuce	20	1/2	Do not apply within 28 days of harvest.
Peanuts	20	1/2	Three applications may be made in a band over the row or broadcast by air or ground equipment. Three applications may be made before digging. One additional application may be made between digging and harvest.
Peppers	20	1/2	Do not apply within 21 days of harvest.
Pumpkins	20	1/2	Do not apply more than once per season nor within 14 days of harvest.
Tablet Beets	30	3/4	Do not apply within 28 days of harvest. Do not eat tops.
Tobacco	20	1/2	Do not apply after plants are 2 feet tall.
Tomatoes	20	1/2	Do not apply within 28 days of harvest. Do not graze treated fields.
Turf and Lawns Ornamentals	20-30	1/2-3/4	Use on ornamental turfs only.

EPA Reg. No. 829-203 A