

Reg # 829-203

Yellow copy

FEB 1 1995

5469878

335
18

Ronald L. Costa Rica
Southern Agricultural Insecticides, Inc.
P.O. Box 218
Palmetto, FL 34220

Dear Mr. Costa Rica:


Subject: SA-50 Insecticide Bait
EPA Registration No. 829-203
Amendment dated 6/1/94 deleting several uses from label

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable provided that you:

1. Make the changes to your label which were required in the Agency letter to you dated June 9, 1994 (enclosed) regarding Worker Protection language.
2. Submit one copy of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions. A stamped copy of the labeling is enclosed for your records. If you have any questions about this letter, please contact Beth Edwards at (703) 305-5400 or (703) 305-6600.

Sincerely yours,



Robert A. Forrest
Product Manager (14)
Insecticide-Rodenticide Branch
Registration Division (7505C)

Enclosures

Edwards

829-203

2/1/95

FEB 1 1995



INSECTICIDE BAIT

CONTAINS 5% DYLOX*

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended for the pesticide registered under EPA Reg. No.

829-203

ACTIVE INGREDIENT:

Dimethyl (2,2,2-trichloro-1-hydroxyethyl) phosphonate*

5.0%

INERT INGREDIENTS:

95.0%

TOTAL

100.0%

*U.S. Patent No. 2,701,225

Trademark of Miles, Inc.

KEEP OUT OF REACH OF CHILDREN

CAUTION

ORGANOPHOSPHATE

May be harmful if swallowed, inhaled, or absorbed through the skin. Avoid contact with water after handling. Do not contaminate feed or food. If poisoning occurs, obtain prompt medical aid. Prolonged exposure will result in cholinesterase depression. **NOTE TO PHYSICIAN:** Atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

This product is toxic to fish, birds, and other wildlife. For terrestrial uses, do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on label. Do not use on other crops used for food or forage. Use only according to label directions. Applications at rates above those shown may result in illegal crop residues. Do not treat food crops grown in the greenhouse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants and shoes plus socks. Follow manufacturers instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker enter into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves, and shoes plus socks.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store pesticide in the original container in a locked storage area inaccessible to children and pets.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke

Follow directions carefully. Timing and method of application, weather conditions and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with the directions and precautions given herewith.

EPA EST No 829 FL-1(3), 829 NC-1(4), 829 NC-2(5) Number following establishment No corresponds to first digit in lot number on bottom of bag

Southern Agricultural Insecticides, Inc.

Palmetto, FL 34220

Hendersonville, NC 28793

Boone, NC 28601

EPA Reg. No 829 203

Net Weight 40 lbs

BEST AVAILABLE COPY

DIRECTIONS FOR USE

Violation of Federal law to use this product in a manner inconsistent with its labeling.

Apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For more information on safety requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Insecticide Bait gives excellent control of surface feeding cutworms, salt marsh caterpillars, armyworms and mole crickets infesting turf and ornamentals. Apply SA-Insecticide Bait as a broadcast treatment over the infested area using a hand or power operated spreader. To help obtain uniform distribution apply one-half of the recommended rate in one direction, then apply the remainder at a right angle.

RECOMMENDED APPLICATIONS

CROP	INSECTS	RATE	REMARKS
ORNAMENTALS Flowers, Shrubs, and Trees	Cutworms	20 - 30 lb. per acre	Obtain thorough coverage.
	Armyworms	(1/2 - 3/4 lb. per 1,000 sq. ft.)	
	Mole Crickets		
TURF and LAWNS*	Cutworms	20 - 30 lb. per acre	Use on ornamental turf only.
	Armyworms	(1/2 - 3/4 lb. per 1,000 sq. ft.)	
	Mole Crickets		

*Do not use treated area or clippings from treated areas for feed or forage.

BEST AVAILABLE COPY