SA 50 BRAND LAWN & TURF INSECTICIDE GRANULES

FOR CONTROL OF: Sod Webworms, White Grubs, Japanese Beetle Grubs, Armyworms, Chinch Bugs, Cutworms, Mole Crickets, Wireworms, Ants, Fleas and Ticks.

ACTIVE INGREDIENTS:


Technical Chlordane Equivalent to 4.2% Octachloro-4,7-methano tetrahydroindane and 2.8% related compounds.)

Carbaryl (1- Naphthyl N-methylcarbamate⁽¹⁾ INERT INGREDIENTS

U.S. Potent No. 2,903,175

For use on ornamental turts only. Do not graze or feed hay from *reated areas

Treat when examination shows prisence of it, etc. or as soon as damage occurs. Ap ply the contents of this bag on 2000 sq. fr. 10 x 50 - utius n. Mour preases will





7.0

7.0% 86.0? 100.0°4

warwaryn (i- iawhannyn ia-mennyncurbamute / INERT INGREDIENTS

U.S. Patent No. 2,903,478

For use on ornamental turfs only. Do not graze or feed hay from *reated areas.

Treat when examination shows presence of insects, or as soon as damage occurs. Apply the contents of this bag on 2,000 sq. ft. (40' x 50') of lawn. Most spreaders will apply the correct amount at the setting for small grass seed, or when using a Cyclone[®] spreader use the No. J setting. Apply to obtain c plete, even coverage. Water turf thoroughly after applying. Examine turf frequently for signs of infestation and re-treat when necessary.

KEEP OUT OF REACH OF CHILDREN

CAUTION: Harmful if swallowed or inhuled. Avoid contact with eyes, skin and clothing. May irritate eyes, nose, throat and skin. Wash thoroughly after handling. Wear clean clothing daily. Store in a safe place. Do not contaminate food or feedstuffs.

Burn empty container. Do not reuse. Stay out of smoke and fumes.

Birds feeding over treated areas may be harmed. This product is toxic to fish and wildlife. Do not contaminate any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. This product is highly toxic to bees exposed to direct or as residues. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Follow instructions carefully. Timing and care of application, weather and lawn conditions, and other factors influencing the results obtained by the use of this product are beyond the control of the seller. Bayer assumes all risks of use, storage, or handling of this material not in strict accordance with the directions and precautions given herewith.

Southern Agricultural Insecticides, Inc.

Palmetto, Fla. FPA Rog. No. 829-206

Handersonville, N. C.

86.0% 100.0%

Borne, N. C.